

## FLASH: A Novel Tool to Identify Vision-Threatening Eye Emergencies

Neil Jairath<sup>1</sup>, Patrick Commiskey<sup>1</sup>, MD; Ariane Kaplan<sup>1</sup>, MD; Yannis M. Paulus<sup>1,2</sup>, MD, FACS

1 Department of Ophthalmology and Visual Sciences, University of Michigan, 1000 Wall Street, Ann Arbor, MI 48105;

2 Department of Biomedical Engineering, University of Michigan, 1000 Wall Street, Ann Arbor, MI 48105.

**Conflict-of-interest statement:** All of the authors have NO affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest; and expert testimony or patent-licensing arrangements), or non-financial interest (such as personal or professional relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed in this manuscript. The authors have no relevant disclosures. All authors have read and contributed substantially to this manuscript and the decision to submit it for publication.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Correspondence to: Yannis M. Paulus, M.D., F.A.C.S., Department of Ophthalmology and Visual Sciences, University of Michigan, Kellogg Eye Center, 1000 Wall Street, Ann Arbor, MI 48105.

Email: [ypaulus@med.umich.edu](mailto:ypaulus@med.umich.edu)

Telephone: +(734) 764-4182

Telephone: +(734) 936-3815

Received: May 19, 2020

Revised: September 10, 2020

Accepted: September 15, 2020

Published online: October 8, 2020

### ABSTRACT

**BACKGROUND:** Two million patients visit emergency departments due to eye complaints annually in the United States, yet nearly one-quarter of these visits are for non-urgent ocular problems. Other

patients often present a significant length of time after the onset of their symptoms, which may cause progression to irreversible vision loss. A major reason for this discrepancy is that many patients are unsure what symptoms constitute eye emergencies. The challenge is helping patients understand what constitutes a vision-threatening eye emergency, as well as the risks and complications that are associated with delaying their visit to the ophthalmologist or Emergency Department.

**OBJECTIVES:** To describe relevant literature on incidence, prevalence, presentation times, associated prognoses, risks, and complications of individual vision-threatening eye emergencies, and present a novel acronym, FLASH (Floaters and flashes, Loss of vision, Aching pain, Second Image, Help), to better educate patients at risk for these conditions, fostering better symptom recognition and timely care. This manuscript is aimed at reaching public health departments, educational institutions, primary care offices and eye care centers as part of a dedicated patient education effort for vision-threatening eye emergencies.

**DESIGN / METHODS:** Narrative overview of the available literature on specific eye conditions presenting with the aforementioned symptoms, synthesizing findings retrieved from searches of computerized databases and authoritative texts.

**CONCLUSIONS:** In each condition presented in this article, symptom interval significantly impacts treatment prognoses. The cited literature demonstrates that patients often present late in emergent eye conditions resulting in vision loss.

**Key words:** Delayed presentation, eye emergencies, vision complications, blindness, patient education, endophthalmitis, retinal detachment

© 2020 The Author(s). Published by ACT Publishing Group Ltd. All rights reserved.

Jairath N, Commiskey P, Kaplan A, Paulus YM. FLASH: A Novel Tool to Identify Vision-Threatening Eye Emergencies. *International Journal of Ophthalmic Research* 2020; 6(1): 336-343 Available from: URL: <http://www.ghrnet.org/index.php/ijor/article/view/2999>

**Financial Support:** This work was supported by the National Eye Institute grant IK08EY027458 (YMP) and the University of Michigan Department of Ophthalmology and Visual Sciences.

## INTRODUCTION

Two million people present every year to emergency departments (ED) for eye complaints in the United States<sup>[1]</sup>. If medical care is not initiated promptly, emergent ophthalmic disease can progress to irreversible vision loss and blindness. However, many of these visits are for non-urgent conditions. A recent study out of the University of Michigan demonstrated that a quarter of eye-related emergency department visits by adults with private insurance over a 14-year period were for non-urgent conditions<sup>[2]</sup>. It is difficult for patients to predict what is and what is not an eye emergency, judging by their ED visits and associated urgency (Table 1). Patients spend nearly \$580 more per visit by presenting to the ED for their ophthalmic concerns compared to an office visit<sup>[2]</sup>. Notably, patients were less likely to seek immediate care for their non-urgent conditions if there were already seeing an eye specialist regularly<sup>[3]</sup>. Public education on vision-threatening entities could foster earlier presentation to eye specialists, cost savings, and maximization of visual prognosis, as well as allowing EDs to serve patients with more appropriate, urgent conditions<sup>[4,5]</sup>.

Many triage tools currently exist and are employed by various eye-care centers around the world. Our team has developed an acronym, described in Figure 1, FLASH (Floaters, Loss of Vision, Aching Pain, Second Vision, Help), that covers the several conditions assessed in these triage tools. Several authors describe examples of triage forms that ask many of the questions FLASH would ask; specifically about double vision, floaters, decreased vision, and pain<sup>[6,7]</sup>. Many academic centers, including the Wills Eye Hospital, include the symptoms listed in FLASH, notably Flashes, Floaters, Sudden Vision Loss, Double

Vision, and Pain as causes for concern for a medical emergency<sup>[7,8]</sup>.

In the future, an increasing number of ophthalmic consultations will likely be performed remotely as telemedicine plays a larger role in our care of patients, causing requests for ophthalmologic consultations to rise rapidly, thus improving eye access for people with emergency conditions<sup>[9]</sup>. With a tool like FLASH, patients would be able to quickly identify and address eye emergencies through such services.

**F** **FLOATERS & FLASHES:**

- New **flashes of light** in the side of your vision, especially if moon-shaped and **increasing** in frequency.
- **Many new** gray or black floaters, or a recent **eye injury**.

**L** **LOSS OF VISION:**

- **Graying or blackening** of vision,
- *even if it gets better* on its own,
- *even if you only lose part* of your vision,
- in either **one or both** eyes.

**A** **ACHING PAIN:**

- **New aching, red eye** and/or **severe light sensitivity**,
- especially if you've had a recent **eye injury**, recent **eye surgery** or are a **contact lens** wearer.

**S** **SECOND IMAGE:**

- **New double vision** that **goes away** by closing **one eye** at a time.

**H** **HELP!**

If you are having these symptoms, please call an ophthalmologist.

- An Ophthalmologist is a physician who specializes in the medical and surgical treatment of eye conditions. Find a near-by Ophthalmologist you. ☎.

**Figure 1** This figure displays the FLASH acronym and its associated symptoms.

**Table 1** This table represents number of people presenting to the emergency department with eye conditions relative to total ED visits, along with demographic breakdown.

	ED visits related to eye injuries*	Treat and release ED visits related to eye injuries*	ED visits resulting in admission related to eye injuries*	ED visits for all other injuries
Number of Visits	636,619	616,766	19,853	29,482,454
ED visits per day	1,744	1,690	54	80,774
Rate per 100,000 population	209	203	7	9,696
Males, rate per 100,000 population	262	254	8	10,400
Females, rate per 100,000 population	158	153	5	9,007
<b>Utilization characteristics</b>				
Percentage treated and released	96.90%			
Percentage admitted to the hospital	3.10%			8.10%
Percentage of visits primarily/principally for eye injuries	80.80%	82.50%	27.60%	n/a
Percentage of visits with a secondary diagnosis of eye injuries only	19.20%	17.50%	72.40%	n/a
<b>Patient Characteristics</b>				
Mean age, years	31.50	30.80	51.20	35.40
<b>Percentage by age group</b>				
0 to 17 years	28.30%	28.80%	12.70%	25.80%
18 to 44 years	45.30%	45.90%	29.10%	41.50%
45 to 64 years	18.70%	18.70%	20.70%	19.20%
65 years and older	7.60%	6.70%	37.60%	13.50%
Percentage of male patients	61.70%	61.80%	59.10%	52.90%
<b>Patient residence, rate per 100,000 population<sup>1</sup></b>				
Urban areas (large central, large fringe, and small and medium metropolitan), rate per 100,000 population	120	116	5	5,317
Rural areas (micropolitan and noncore), rate per 100,000 population	646	630	16	31,229

<sup>1</sup>Missing patient's residence on 250,418 (<1%) injury records and income information on 751,631 (2.5%) injury records. \*Based on all-listed diagnoses. Source: Agency for Healthcare Research and Quality, Healthcare Cost and Utilization Project, Nationwide Emergency Department Sample, 2008. Denominator data for rates were based on Annual Estimates of Resident Population for the United States, Regions, States and Puerto Rico: April 1, 2000 to July 1, 2009 (NST\_EST2009-01, NC-EST2009-02). U.S. Census Bureau, Population Division.

Clinical presentation	Score 1	Score 2	Score 3	Score 4
Marked decrease in best visual acuity (worse or = 20/60), or 4 lines decrease compared to last visual assessment.	Shaded	Shaded	Shaded	Unshaded
Mild decrease in best visual acuity (better than 20/60), or 2 lines decrease compared to last visual assessment.	Unshaded	Shaded	Shaded	Shaded
Peripheral loss of vision	Shaded	Shaded	Shaded	Unshaded
IOP more than 25 mm Hg by air puff	Shaded	Unshaded	Shaded	Shaded
Mild ocular pain*/foreign body sensation	Shaded	Shaded	Shaded	Shaded
Severe ocular pain*	Shaded	Unshaded	Shaded	Shaded
Mild eye/eyelid redness†	Shaded	Shaded	Shaded	Shaded
Severe eye/eyelid redness†	Shaded	Unshaded	Shaded	Shaded
Eye redness in a preschool age	Shaded	Shaded	Shaded	Unshaded
Contact lens wearer	Shaded	Shaded	Shaded	Shaded
History of corneal transplant	Shaded	Shaded	Shaded	Shaded
Proptosis	Shaded	Unshaded	Shaded	Shaded
New onset ptosis	Shaded	Shaded	Shaded	Shaded
Double vision	Shaded	Shaded	Shaded	Shaded
Double vision with pupil asymmetry	Shaded	Shaded	Shaded	Unshaded
Flashes	Shaded	Unshaded	Shaded	Shaded
Floater	Shaded	Shaded	Shaded	Shaded
Suspect traumatic open globe injury	Shaded	Shaded	Unshaded	Shaded
Suspect chemical burn	Shaded	Shaded	Shaded	Unshaded
Child less than 6 months	Shaded	Shaded	Shaded	Shaded
<b>Total Score</b>				
Time of registration: _____ Time of screening: _____ Time of examination: _____				
Classification: Non ER (total score of 1 or less)		ER (total score of 2 and 3)		Top ER (total score 4 or more)
Final diagnosis (possible diagnosis): _____				
Name/Signature ER Nurse: _____ Name /Signature ER Resident: _____				

\*mild ocular pain means pain not preventing the patient carrying out usual activities such as sleeping and routine work. Severe pain prevents the patient from carrying out usual activities, †mild/severe eye redness will be scored based on eye photo posted in emergency room (ER). Unshaded boxes denotes the allowed space for entry. IOP - intraocular pressure

**Figure 2** This figure is obtained from reference [6] and represents the clinical presentation of emergency ophthalmic patients. The score delineates the urgency of the condition: non-urgent (0 to 1), semi-urgent (2 or 3), and urgent (4 or more). The FLASH acronym includes most of the symptoms that would be recognizable to a patient experiencing them. Copyright: © Saudi Medical Journal.

In this manuscript, we have undertaken a review of the relevant literature on incidence, prevalence, presentation times, associated prognoses, risks, and complications of individual vision-threatening eye emergencies that present with the symptoms outlined in FLASH. We hope to demonstrate the need for patient education on these emergencies, as far too many patients are currently suffering from preventable complications.

**2.1: Retinal detachment (Floaters / Flashes, Loss of Vision)**

Macula-sparing rhegmatogenous retinal detachments (RRD) are urgent eye conditions in which the neurosensory retina is detached from the underlying retinal pigment epithelium and choroid, which if left untreated, can result in permanent blindness. RRD has an incidence of about 5 cases in 100,000 persons per year (for otherwise normal eyes), but 20 in 100,000 when including only middle-age or elderly populations<sup>[10,11]</sup>. Lifetime risk for RRD is approximately 1 in 300, and notably shows a strong correlation with affluence, indicating that RRD may become more prevalent as populations age in more affluent countries<sup>[12,13]</sup>.

In the case of retinal detachment, patients often present once vision declines and the macula is off, leading to a worse visual prognosis and an average visual acuity outcome of 20/80. If caught earlier when the macula is on, the prognosis is much better<sup>[14]</sup>. Estimates have determined that between 50% to 70% of patients (regardless of their level of education) present after the macula is detached, due to

their not being able to recognize the symptoms of detachment<sup>[15,16]</sup>, but those who present with macula still on are more sensitive to the presence of floaters and seem to be more informed about signs and symptoms of eye conditions. This awareness is essential to achieving earlier presentation times and can become more widespread with promotion strategies such as educating patients about warning signs (especially in high risk patients, such as high myopes and the elderly)<sup>[17,18]</sup>.

One study showed that length of delay in surgical intervention from onset of symptoms correlated strongly with a decline in visual acuity, with 53% achieving 20/20 to 20/50 acuity within 9 days, 34% between 10-19 days, and 29% after 19 days. Patients who presented within 9 days were significantly more likely to obtain a final acuity of 20/50 or better ( $p < 0.05$ )<sup>[19]</sup>. Another study demonstrated that after detachment, the best mean preoperative vision, the most significant predictor of postoperative visual acuity<sup>[20]</sup>, is seen in patients with detachment of less than 1 week duration. If the macula has been off for longer than 6 weeks, visual prognosis postoperatively is significantly poorer<sup>[21,22]</sup>. In order to prevent the complication of blindness from this treatable condition, early diagnosis and prompt referral for diagnosis and surgery are of paramount importance.

**2.2 Acute angle closure glaucoma (Loss of vision, Acute pain)**

Acute angle closure glaucoma (AACG) is an eye emergency in which closure of the drainage system of the eye, the angle, causes an

acute elevation in intraocular pressure (IOP) and can lead to ischemia and permanent vision loss if untreated. Awareness and knowledge of glaucoma is low in patients suffering from AACG. This lack of awareness, leading to delayed presentation times, is correlated with many factors, including older age, education, and unemployment<sup>[23]</sup>. Most symptomatic angle closure events stand a high chance of recovery without any short-term optic disc or visual field damage if promptly treated (60-75% success rate)<sup>[24]</sup>.

Development of primary angle closure glaucoma after AACG was noted in two studies of Caucasian and Asian populations to correlate with length of symptoms before presentation ( $p = 0.001$ ) and time taken to abort the acute attack ( $p = 0.01$ )<sup>[25,26]</sup>. In a Scottish population, delayed presentation ( $\geq 3$  days) was associated with higher rate of glaucoma at follow-up (22.6% vs 60.8%,  $p < 0.001$ ), worse visual acuity (20/44 vs 20/110 Snellen,  $p < 0.0001$ ) and need for more topical medication (RR 0.52 vs 1.2,  $p = 0.003$ ) to control IOP<sup>[27]</sup>.

Visual field loss has been shown to be directly correlated to delayed presentation after an AACG attack, with one study demonstrating loss of vision in 38% of eyes after nerve fiber loss six months after the attack<sup>[28]</sup>. Another study demonstrated that changes in optic disc morphology were documented as soon as 2-16 weeks after the attack, preferentially targeting inferotemporal and superotemporal regions<sup>[28]</sup>. Although IOP measurement is an important predictor of outcome, if an attack persists for a longer duration and becomes more difficult to manage, the visual outcome for the eye worsens, regardless of initial IOP measurement<sup>[29,30]</sup>. One study assessed the risk of chronic glaucoma in patients who presented with a delay as well as patients with poor IOP control, and found that the relative risk of patients presenting with a 24 to 72-hour delay was 2.78, while patients requiring iridotomy or trabeculectomy for control of IOP had relative risks of 3.63 and 4.83, respectively<sup>[30]</sup>.

### 2.3 Corneal Ulcer (Loss of vision, Acute Pain)

Corneal ulcers are infections, most commonly bacterial, in the cornea which can lead to permanent scarring and visual loss. The surge in use of contact lenses has led to an increasing prevalence of corneal ulcers in recent years.<sup>31</sup> Incidence of microbial keratitis in the population varies based on type of contact lens used, but is clearly correlated to length of time in the eye, with yearly rates of 2 cases per 10,000 people for rigid contact lens wearers, 2.2-4.4 per 10,000 for daily-wear soft contact lens users, and 13.3-20.9 per 10,000 for extended-wear soft contact lenses<sup>[32]</sup>. A known complication of corneal ulcers is permanent blindness, and a delay in diagnosis and treatment can lead to this unfortunate result<sup>[33]</sup>. For this reason, immediate referral to the ophthalmologist, leading to prompt initiation of antimicrobial therapy, is paramount to preventing loss of vision in these patients.

Increased rates of recurrence and necessity for invasive operations are seen in patients with delayed presentation<sup>[34]</sup>. The average time before a hospital visit after onset of symptoms in Nadu, India, was 6.2 days, and 10.6% of cases either required therapeutic penetrating keratoplasty (8.5%) or progressed to endophthalmitis and required evisceration (2.1%)<sup>[34]</sup>. In studies out of Iran and Britain, the average delay between onset of symptoms and the first examination was 48 hours, and complications requiring intervention beyond a topical approach were noted in 6% of eyes<sup>[35,36]</sup>. In one study, the mean delay in referral to the corneal specialist for patients that had to undergo penetrating keratoplasty was 8 days, versus 1.3 days in the patients that did not ( $p < 0.03$ )<sup>[37]</sup>.

Delayed presentation may negate the efficacy of medical treatment, according to authors who found that when *Fusarium* keratitis has progressed to an advanced stage, combination antibiotic therapies, in-

cluding fluconazole, ketoconazole, topical natamycin, and amphotericin B failed to elicit a response in 9 of 10 patient cases<sup>[38]</sup>. A symptom interval of more than 2 weeks in patients with fungal keratitis has also been shown to statistically increase the chances that they will require surgery later<sup>[39]</sup>. However, if diagnostic delay and initiation of antibiotic therapy can be held to under 18 days from the start of symptoms, the odds of a better final visual acuity are increased<sup>[40]</sup>. For these reasons, early presentation and diagnosis are important factors in determining prognosis for visual outcome in patients with corneal ulcers.

### 2.4 Herpes Keratitis (Loss of vision, Acute pain)

Herpes can cause a particularly devastating corneal infection which can lead to scarring, vision loss, and blindness. The incidence of ocular herpes simplex virus (HSV) infection in the USA is about 0.15%, with about 20,000 new cases and 28,000 reactivations annually. In the United States, ocular HSV is widely known to be a frequent cause of blindness<sup>[41]</sup>. Some estimations predict that nearly 500,000 people are afflicted with ocular HSV in the US<sup>[42]</sup>. Necrotizing stromal keratitis (NSK), a result of the progression of ocular HSV, is a condition that can lead to corneal perforation and vision loss. The diagnosis of NSK is often delayed, leading to increased incidence of visual morbidity<sup>[43]</sup>.

In most cases, ocular HSV will resolve on its own without permanently damaging vision, with mean times to resolution after symptom onset ranging from 17.6 days in the first episode to 28.4 days in recurrent episodes<sup>[41]</sup>. Notably, there is no current method that will predict which infections may progress, so prompt recognition and treatment is important in preventing progression, and alleviating symptoms<sup>[44]</sup>. Despite its often self-limited nature, reducing viral replication is imperative to reduce symptoms, shorten the course of disease, and prevent further complications<sup>[45]</sup>.

### 2.5 Scleritis (Loss of vision, Acute pain)

Scleritis is a painful inflammation of the sclera and scleral vessels that is associated with systemic autoimmune conditions in up to 50% of patients. Scleritis is highly variable in its presentation, with time from an initial identifiable event to presentation ranging anywhere from 0-36 years. Postsurgical patients tend to fall on the longer end of the spectrum compared to post-nonsurgical trauma patients<sup>[46]</sup>. Autoimmune disease processes are the most common etiology for scleritis and can be progressively destructive if treatment is not initiated, potentially leading to loss of vision or loss of the eye<sup>[47]</sup>. Fungal etiologies tend to have worse prognosis, likely secondary to delays in diagnosis<sup>[48]</sup>. Infectious scleritis can be viral, bacterial, fungal, and parasitic. It is uncommon, particularly in the absence of infectious keratitis; however, possibly because of the diagnostic delay that is reported to be associated with infectious scleritis, visual prognosis is poorer in these patients than in those with an autoimmune manifestation<sup>[49]</sup>. One study reported that there is a gap of nearly 46 days between commencing treatment and resolution of scleritis, which can be exacerbated by a delayed diagnosis: Hodson *et al.* report 50% of eyes losing function vision, emphasizing the point that delayed diagnosis can result in progression of the disease that can result in eventual vision loss<sup>[46]</sup>.

### 2.6 Giant cell arteritis (Loss of vision)

Giant cell arteritis is the most common vasculitis in adults aged 50 years and older and can often present with eye complaints. The overall prevalence of giant cell arteritis (GCA), adjusted for age and sex, has been reported to be nearly 204 (95% CI: 161-254) per 100,000

population greater than or equal to 50 years of age, with a predilection for men<sup>[50]</sup>. Nearly 15-20% of patients with GCA have reported experiencing permanent vision loss as a complication of this condition, which makes early and correct diagnosis critical<sup>[51,52]</sup>. During the course of disease progression, nearly 50% of patients experience visual symptoms, including painless unilateral vision blurring or vision loss and diplopia. What initially seems like an innocuous partial field defect can progress to blindness over the course of a few days. Although transient episodes of blurred vision are generally treatable, if treatment for GCA is not started urgently, sudden irreversible vision loss can result. One of the most important considerations in treatment of the GCA is the potential for bilateral vision loss. If GCA remains untreated in patients with unilateral vision loss, the second eye may become affected within 1-2 weeks in up to 50% of patients<sup>[53]</sup>.

In the classic disease progression of GCA, it tends to involve the aorta and collateral branches, which can lead to a host of complications. Besides a permanent loss of visual function, untreated GCA, potentially due to late diagnosis, can cause serious systemic complications<sup>[52]</sup>. And delay in diagnosis is not uncommon, with a reported mean delay of 9 weeks, and even longer when patients displayed an absence of cranial symptoms<sup>[54]</sup>. In one study, the mean time from symptom onset to diagnosis of GCA was 35 days due to slow recognition of ischemic symptoms. In the 65-patient cohort, 24.6% experienced visual loss at presentation, 15.4% presented with neuro-ophthalmological vascular complications (NOC), 70% of whom developed permanent visual impairment, and 7.7% suffered cerebrovascular complications. A delay in diagnosis and/or treatment has been found to partially contribute to the ischemic manifestations that result from GCA, most of which are not reversible<sup>[55]</sup>.

### 2.7 Endophthalmitis (Loss of vision, Acute pain)

Endophthalmitis is a severe inflammation of the intraocular tissues, most commonly due to infection. It is particularly devastating since the eye, like the brain, is an immune-privileged site, and thus inflammation can lead to permanent, severe vision loss. Because symptoms of endophthalmitis (both systemic and ocular) are generally non-specific, diagnosis is reliant on patients contacting clinicians with their symptoms. With continuing progress and advancement in surgical technology, as well as vitreoretinal pharmacology, prompt intervention with intravitreal pharmacotherapy or surgery (in more severe cases) is the prevailing trend in endophthalmitis management<sup>[56]</sup>. Endophthalmitis may progress into permanent vision loss, and this can happen in a relatively short time window<sup>[57]</sup>.

Endophthalmitis generally presents within a few hours to a few days after the onset of symptoms. Cases of endophthalmitis are considered medical emergencies. Despite the risk of blindness, vision may be preserved if prompt therapy is initiated, even in the most severe cases<sup>[58]</sup>. Most patients suffering from postoperative endophthalmitis present within seven days of the surgery with an acute onset<sup>[59]</sup>. Acute bacterial endophthalmitis is an especially emergent condition, as even a short treatment delay can lead to loss of vision<sup>[60]</sup>. A 19-year-long study out of the Bascolm Palmer Eye Institute found that for cases occurring within six weeks of clear corneal cataract surgery, time to diagnosis ranged from 1-39 days, with a median of 9 days and a mean of 13.8<sup>[61]</sup>. It has been widely reported that longer delay from infection to treatment correlates directly to a poorer visual prognosis<sup>[62-64]</sup>. One study examining retinal changes in the setting of *Bacillus* endophthalmitis demonstrated structural and electrophysiologic changes, PMN infiltration into the vitreous, Glial fibrillary acidic protein (GFAP) detection in the retina, and an increase in ocular TNF- $\alpha$ , all within 4 to 6 hours. The changes in retinal

architecture correlated to a functional deficit in these eyes. The study suggests that prompt intervention in endophthalmitis is strongly indicated, due to the rapid detrimental changes that can occur to the eye, potentially leading to vision loss, in patients with endophthalmitis<sup>[65]</sup>.

### 2.8 New-onset diplopia (Loss of vision, Second vision)

Although this manuscript has been organized by condition, rather than symptom, the eye emergencies associated with new-onset diplopia are specific and severe enough to be presented as a symptom with associated causes. Specific associated emergency conditions include cranial nerve (CN) III, IV, or VI palsy, cavernous sinus syndrome (CSS), or CNS lesions.

CN palsy, according to a study out of Moorfields Eye Hospital, is the most common condition affecting patients who present to the clinic complaining of diplopia in both eyes, with a prevalence of 67%<sup>[66]</sup>. One cause of a CN palsy may be Cavernous Sinus Syndrome (CSS). Prompt diagnosis and treatment within 100 days of onset of CSS plays an important role in improved visual prognosis<sup>[67,68]</sup>.

Diplopia is also known to present as a symptom of CNS lesions, such as aneurysm rupture or brain tumor. One of the most worrisome complications that can arise from aneurysms that have not yet ruptured is subarachnoid hemorrhage, occurring at a rate of nearly 6-10/100,000 person-years. Rupture of these unruptured intracranial aneurysms (UIAs) are responsible for nearly 85% of all SAH cases, with a fatality rate of 30%-40%<sup>[69,70]</sup>. Visual difficulties (including diplopia) accounted for 38% of symptoms occurring at any time and for 10% of the first presenting symptoms of brain tumor. The average symptom interval, correlated with a negative impact on treatment outcome, for patients presenting with brain tumors was 471 days with a median of 120 days<sup>[71]</sup>. Diplopia was present in 43% of these cases, while blurred vision was present in 39%. Photophobia was present in 6% of cases<sup>[72]</sup>.

In one study out of the UK that examined the records of 139 children with brain tumors, it was found that a delay in diagnosis was related to many complications including reduced visual acuity, growth abnormalities, cranial nerve palsy, and head tilt. The median time to diagnosis from the onset of symptoms was 3.3 months, by which time patients usually had around six signs and/or symptoms<sup>[73]</sup>. Separate studies have corroborated this timeframe, with a reported lag time from symptom onset to diagnosis of at least 12.5 weeks<sup>[74-77]</sup>.

## CONCLUSION

Patients have a limited understanding about what constitutes vision-threatening eye emergencies, and what symptoms to look for when they feel like they may be experiencing one. This lack of information may result in devastating side effects, including progression to irreversible vision loss and blindness. In addition, patients not presenting in a timely fashion to ophthalmic emergencies and seeking emergency department treatment for non-urgent conditions incur significant cost to the healthcare system (estimated \$392.9 million every year)<sup>[3,78]</sup>.

Currently, it is apparent that clinical management of vision-threatening eye emergencies can most effectively help those who seek clinical help for appropriate conditions in a timely fashion. In every condition presented in this article, time to presentation after onset of symptoms was an important factor in determining treatment prognoses. The current challenge, however, is helping patients to understand what constitutes a vision-threatening eye emergency, as well as the risks and complications that are associated with delaying their visit to the ophthalmologist.

Through a narrative review of the literature discussing the delayed presentation of several serious ophthalmic emergencies and the decreased outcomes associated with time to presentation, including symptom recurrence, irreversible vision loss, and blindness, we hope to have demonstrated the need for a public health education initiative that would prompt patients to present to their clinicians earlier after the onset of symptoms and before permanent damage is done. Although in some cases, the literature is sparse, this speaks to the relative obscurity of the subject, and the need to shed light on vision-threatening, and in some cases, life-threatening emergencies. We recognize this as a potential limitation to the study. We propose one step toward a solution in the form of an acronym, much like the widely successful FAST acronym for stroke used as a mnemonic to help detect and enhance responsiveness to stroke victim needs. Through demonstration of the need for a dedicated public health effort to serve this problem, and the introduction of a potentially useful acronym to help achieve this goal, patients will be encouraged to present earlier to physicians if they are experiencing vision-threatening eye emergencies.

The acronym we propose is **FLASH**: **F** stands for Floaters and Flashes, **L** stands for Loss of Vision, **A** stands for Acute Pain, **S** stands for Second Image (double-vision), and **H** stands for Help (contact your ophthalmologist). The medical community has already seen massive success in the implementation of the FAST / BEFAST acronyms for stroke, as patients at risk are more informed about the signs to look for and possess an acronym that is relevant and easy to remember. By introducing FLASH, we hope to implement the same model into primary care offices and eye care centers, with the aim of improving patient outcomes for these emergencies.

## REFERENCES

- Channa R, Zafar SN, Canner JK, Haring RS, Schneider EB, Friedman DS. Epidemiology of eye-related emergency department visits. *JAMA Ophthalmol.* 2016; **134**(3): 312-319. [PMID: 26821577]; [DOI: 10.1001/jamaophthalmol.2015.5778]
- Network for Excellence in Health Innovation (2010). A matter of urgency: reducing emergency department overuse; 2010. Available at: [http://www.nehi.net/writable/publication\\_files/file/nehi\\_ed\\_overuse\\_issue\\_brief\\_032610final.pdf](http://www.nehi.net/writable/publication_files/file/nehi_ed_overuse_issue_brief_032610final.pdf).
- Uhr JH, Mishra K, Wei C, Wu AY. Awareness and Knowledge of Emergent Ophthalmic Disease Among Patients in an Internal Medicine Clinic. *JAMA Ophthalmol.* 2016; **134**(4): 424-31. [PMID: 26892039]; [DOI: 10.1001/jamaophthalmol.2015.6212]
- Graves D. Triaging Ocular Emergencies. Certified Patient Service Specialist, 2012. [PMID: 23270836]; [DOI: 10.1016/j.mayocp.2020.03.018]
- Fleming, TC. Emergency Triage - Protecting the Patients and the Practice. Paraoptometric Resource Center, 2013. American Optometric Association.
- Alsamnan MS, Mousa A, Al-kuwaileet S, Al-Suhaibani AH. Triaging self-referred patients attending ophthalmic emergency room. *Saudi Med J.* 2015; **36**(6): 678-84. [PMID: 25987109]; [DOI: 10.15537/smj.2015.6.11302]
- Wills Eye Hospital. Eye Emergency Department. Available at: [www.willseye.org/eye-emergency-department](http://www.willseye.org/eye-emergency-department). 2013.
- Stagg, BC, Shah MM, Talwar N, Padovani-Claudio DA, Woodward MA, Stein JD. Factors Affecting Visits to the Emergency Department for Urgent and Nonurgent Ocular Conditions. *Ophthalmol.* 2017; **124**(5): 720-729. [PMID: 28159379]; [DOI: 10.1016/j.ophtha.2016.12.039]
- ScienceDaily. Telemedicine could improve eye exam access for people with diabetes. Available at: <https://www.sciencedaily.com/releases/2016/08/160829095606.htm>. 2018.
- Li X, Beijing Rhegmatogenous Retinal Detachment Study Group. Incidence and epidemiological characteristics of rhegmatogenous retinal detachment in Beijing, China. *Ophthalmol.* 2003; **110**(12): 2413-2417. [PMID: 14644727]; [DOI: 10.1016/s0161-6420(03)00867-4]
- Mitry D, Charteris DG, Fleck BW, Campbell H, Singh J. The epidemiology of rhegmatogenous retinal detachment: geographical variation and clinical associations. *Br J Ophthalmol.* 2010; **94**(6): 678-84. [PMID: 19515646]; [DOI: 10.1136/bjo.2009.157727]
- Gariano RF, Kim CH. Evaluation and management of suspected retinal detachment. *Am Fam Physician.* 2004; **69**(7): 1691-8. [PMID: 15086041]
- Mitry D, Charteris DG, Yorston D, Rehman Siddiqui, MA, Campbell H, Murphy AL, Fleck BW, Wright AF, Singh J, Scottish RD Study Group. The epidemiology and socioeconomic associations of retinal detachment in Scotland: a two-year prospective population-based study. *Invest Ophthalmol Vis Sci.* 2010; **51**(10): 4963-8. [PMID: 20554615]; [DOI: 10.1167/iov.10-5400]
- Doyle E, Herbert EN, Bunce C, Williamson TH, Laidlaw DAH. How effective is macula-off retinal detachment surgery. Might good outcome be predicted?. *Eye.* 2007; **21**(4): 534-40. [PMID: 16456590]; [DOI: 10.1038/sj.eye.6702260]
- Quintyn JC, Benouaich X, Pagot-mathis V, Mathis A. Retinal detachment, a condition little known to patients. *Retina.* 2006; **26**(9): 1077-8. [PMID: 17151498]; [DOI: 10.1097/01.iae.0000254889.71142.ce]
- Goezinne F, La heij EC, Berendschot TT, Tahzib NG, Koetsier LS, Hoevenaars JGMM, Liem ATA, Kijlstra A, Webers CAB, Hendrikse F. Patient ignorance is the main reason for treatment delay in primary rhegmatogenous retinal detachment in The Netherlands. *Eye.* 2009; **23**(6): 1393-9. [PMID: 18791551]; [DOI: 10.1038/eye.2008.272]
- Eijk ES, Busschbach JJ, Timman R, Monteban HC, Vissers JMH, van Meurs JC. What made you wait so long? Delays in presentation of retinal detachment: knowledge is related to an attached macula. *Acta Ophthalmol.* 2016; **94**(5): 434-40. [PMID: 27008986]; [DOI: 10.1111/aos.13016]
- Rehman Siddiqui MA, Abdelkader E, Hammam T, Murdoch JR, Lois N. Socioeconomic status and delayed presentation in rhegmatogenous retinal detachment. *Acta Ophthalmol.* 2010; **88**(8): e352-3. [PMID: 19900198]; [DOI: 10.1111/j.1755-3768.2009.01771.x]
- Burton TC. Recovery of visual acuity after retinal detachment involving the macula. *Trans Am Ophthalmol Soc.* 1982; **80**: 475. [PMID: 6763802]
- Sharma T, Challa JK, Ravishankar KV, Murugesan R. Scleral buckling for retinal detachment. Predictors for anatomic failure. *Retina.* 1994; **14**: 338. [PMID: 7817027]; [DOI: 10.1097/00006982-199414040-00008]
- Mowatt L, Shun-shin GA, Arora S, Price N. Macula off retinal detachments. How long can they wait before it is too late?. *Eur J Ophthalmol.* 2005; **15**(1): 109-17. [PMID: 15751248]
- Kang HK, Luff AJ. Management of retinal detachment: a guide for non-ophthalmologists. *BMJ.* 2008; **336**(7655): 1235-40. [PMID: 18511798]; [DOI: 10.1136/bmj.39581.525532.47]
- Saw SM, Gazzard G, Friedman D, Foster PJ, Devereux JG, Wong, ML, Seah S. Awareness of glaucoma, and health beliefs of patients suffering primary acute angle closure. *Br J Ophthalmol.* 2003; **87**(4): 446-9. [PMID: 12642308]; [DOI: 10.1136/bjo.87.4.446]
- Douglas GR, Drance SM, Schulzer M. The visual field and nerve head in angle-closure glaucoma. A comparison of the effects of acute and chronic angle closure. *Arch Ophthalmol.* 1975; **93**: 409-411. [PMID: 1131080]; [DOI: 10.1001/archophth.1975.01010020423004]
- Andreatta W, Elaroud I, Nightingale P, Nessim M. Long-term outcomes after acute primary angle closure in a White Caucasian

- population. *BMC Ophthalmol.* 2015; **15**: 108. [PMID: 26286533]; [DOI: 10.1186/s12886-015-0100-5]
26. Tan AM, Loon SC, Chew PT. Outcomes following acute primary angle closure in an Asian population. *J Clin Exp Ophthalmol.* 2009; **37(5)**: 467-72. [PMID: 19624342]; [DOI: 10.1111/j.1442-9071.2009.02060.x]
  27. Chua PY, Day AC, Lai KL, Hall N, Tan LL, Khan K, Lim LT, Foot B, Foster PJ, Azuara-Blanco, A. The incidence of acute angle closure in Scotland: a prospective surveillance study. *Br J Ophthalmol.* 2017. [PMID: 28794074]; [DOI: 10.1136/bjophthalmol-2017-310725]
  28. Shen SY, Baskaran M, Fong AC, Chan YH, Lim LS, Gazzard, G, Seah, SKL, Aung, T. Changes in the optic disc after acute primary angle closure. *Ophthalmol.* 2006; **113**: 924-929. [PMID: 16751035]; [DOI: 10.1016/j.ophtha.2006.01.070]
  29. David R, Tessler Z, Yassur Y. Long-term outcome of primary acute angle-closure glaucoma. *Br J Ophthalmol.* 1985; **69**: 261-262. [PMID: 3994941]; [DOI: 10.1136/bjo.69.4.261]
  30. Wong JS, Chew PT, Alsagoff Z, Poh K. Clinical course and outcome of primary acute angle-closure glaucoma in Singapore. *Singapore Med J.* 1997; **38**: 16-18. [PMID: 9269347]
  31. Liesegang TJ. Contact lens-related microbial keratitis: Part I: Epidemiology. *Cornea.* 1997 Mar; **16(2)**: 125-31. [PMID: 9071523]
  32. Miedziak AI, Miller MR, Rapuano CJ, Laibson LR, Cohen EJ. Risk factors in microbial keratitis leading to penetrating keratoplasty. *Ophthalmol.* 1999; **106(6)**: 1166-70.
  33. Loh K, Agarwal P. Contact lens related corneal ulcer. *Malays Fam Physician.* 2010; **5(1)**: 6-8.
  34. Butler TK, Males JJ, Robinson LP, Wechsler AW, Suttonm GL, Cheng J, Taylor P, McClellan K. Six-year review of Acanthamoeba keratitis in New South Wales, Australia: 1997-2002. *J Clin Exp Ophthalmol.* 2005; **33(1)**: 41-6. [PMID: 10366087]; [DOI: 10.1016/S0161-6420(99)90250-6]
  35. Rathi H, Venugopal A, Rameshkumar G, Ramarkishnan R, Meenakshi R. Fungal Keratitis Caused by Exserohilum, An Emerging Pathogen. *Cornea.* 2016; **35(5)**: 644-646. [PMID: 26863500]; [DOI: 10.1097/ico.0000000000000774]
  36. Hedayati H, Ghaderpanah M, Rasoulinejad SA, Montazeri M. Clinical Presentation and Antibiotic Susceptibility of Contact Lens Associated Microbial Keratitis. *J Pathog.* 2015; **2015**: 152767. [PMID: 26770828]; [DOI: 10.1155/2015/152767]
  37. Bourcier T, Thomas F, Borderie V, Chaumeil C, Laroche L. Bacterial keratitis: predisposing factors, clinical and microbiological review of 300 cases. *Br J Ophthalmol.* 2003; **87(7)**: 834-8. [PMID: 12812878]; [DOI: 10.1136/bjo.87.7.834]
  38. Vajpayee RB, Dada T, Saxena R, Vajpayee, M, Taylor HR, Venkatesh P, Sharma N. Study of the first contact management profile of cases of infectious keratitis: a hospital-based study. *Cornea.* 2000; **19**: 52-6. [PMID: 10632009]; [DOI: 10.1097/00003226-200001000-00011]
  39. Dursun D, Fernandez V, Miller D, Alfonso EC. Advanced fusarium keratitis progressing to endophthalmitis. *Cornea.* 2003 May; **22(4)**: 300-3. [PMID: 12792470]; [DOI: 10.1097/00003226-200305000-00004]
  40. Keay LJ, Gower EW, Iovieno A, Oechsler RA, Alfonso EC, Matoba A, Colby K, Tuli SS, Hammersmit K, Cavanagh D, Lee SM, Irvine J, Stulting RD, Mauger TF, Schein OD. Clinical and microbiological characteristics of fungal keratitis in the United States, 2001-2007: a multicenter study. *Ophthalmol.* 2011; **118(5)**: 920-6. [PMID: 21295857]; [DOI: 10.1016/j.ophtha.2010.09.011]
  41. Liesegang TJ, Melton LJ 3rd, Daly PJ, Ilstrup DM. Epidemiology of ocular herpes simplex. Incidence in Rochester, Minn, 1950 through 1982. *Arch Ophthalmol.* 1989 Aug. **107(8)**: 1155-9. [PMID: 2787981]; [DOI: 10.1001/archophth.1989.01070020221029]
  42. Lairson DR, Begley CE, Reynolds TF, Wilhelmus KR. Prevention of herpes simplex virus eye disease: A cost-effectiveness analysis. *Arch Ophthalmol.* 2003; **121**: 108-12. [PMID: 12523894]; [DOI: 10.1001/archophth.121.1.108]
  43. Knickelbein JE, Hendricks RL, Charukamnoetkanok P. Management of herpes simplex virus stromal keratitis: An evidence-based review. *Surv Ophthalmol.* 2009; **54**: 226-34. [PMID: 19298901]; [DOI: 10.1016/j.survophthal.2008.12.004]
  44. McDonald, M, Hardten DR, Mah FS, O'Brien TP, Rapuano CJ, Schanzlin DJ, Shamie N, Sheppard JD, Tauber S, Waring GO. Management of Epithelial Herpetic Keratitis: An Evidence-Based Algorithm. Report, Bausch and Lomb, Chicago, 2012
  45. Azher TN, Yin XT, Tajfirouz D, Huang AJ, Stuart PM. Herpes simplex keratitis: challenges in diagnosis and clinical management. *Clin Ophthalmol.* 2017; **11**: 185-191. [PMID: 28176902]; [DOI: 10.2147/OPHTH.S80475]
  46. Hodson KL, Galor A, Karp CL, Davis JL, Albin TA, Perez VL, Miller D, Forster RK. Epidemiology and visual outcomes in patients with infectious scleritis. *Cornea.* 2013 Apr; **32(4)**: 466-72. [PMID: 22902495]; [DOI: 10.1097/ICO.0b013e318259e952]
  47. Jabs DA, Mudunn A, Dunn JP, Marsh MJ. Episcleritis and scleritis: clinical features and treatment results. *Am J Ophthalmol.* 2000; **130**: 469-476. [PMID: 11024419]; [DOI: 10.1016/S0002-9394(00)00710-8]
  48. Ahn SJ, Oh JY, Kim MK, Lee JH, Wee WR. Clinical features, predisposing factors, and treatment outcomes of scleritis in the Korean population. *Korean J Ophthalmol.* 2010 Dec; **24(6)**: 331-5. [PMID: 21165230]; [DOI: 10.3341/kjo.2010.24.6.331]
  49. Rameneden R, Raiji V. Clinical characteristics and visual outcomes in infectious scleritis: a review. *Clin Ophthalmol.* 2013; **7**: 2113-22. [PMID: 24235809]; [DOI: 10.2147/OPHTH.S37809]
  50. Crowson CS, Matteson EL. Contemporary prevalence estimates for giant cell arteritis and polymyalgia rheumatica, 2015. *Semin Arthritis Rheum.* 2017; **47(2)**: 253-256. [PMID: 28551169]; [DOI: 10.1016/j.semarthrit.2017.04.001]
  51. Aiello PD, Trautmann JC, McPhee TJ, Kunselman AR, Hunder GG. Visual prognosis in giant cell arteritis. *Ophthalmol.* 1993; **100(4)**: 550-5. [PMID: 8479714]; [DOI: 10.1016/s0161-6420(93)31608-8]
  52. Soriano A, Muratore F, Pipitone N, Boiardi L, Cimino L, Salvareni C. Visual loss and other cranial ischaemic complications in giant cell arteritis. *Nat Review Rheumatol.* 2017; **13(8)**: 476-484. [PMID: 28680132]; [DOI: 10.1038/nrrheum.2017.98]
  53. Liozon E, Ly KH, Robert PY. [Ocular complications of giant cell arteritis]. *Rev Med Interne.* 2013; **34(7)**: 421-30. [PMID: 23523078]; [DOI: 10.1016/j.revmed.2013.02.030]
  54. Prior JA, Ranjbar H, Belcher J, Mackie SL, Helliwell T, Liddle J, Mallen CD. Diagnostic delay for giant cell arteritis - a systematic review and meta-analysis. *BMC Med.* 2017; **15(1)**: 120. [PMID: 28655311]; [DOI: 10.1186/s12916-017-0871-z]
  55. Ezeonyeji AN, Borg FA, Dasgupta B. Delays in recognition and management of giant cell arteritis: results from a retrospective audit. *Clin Rheumatol.* 2011; **30(2)**: 259-62. [PMID: 21086005]; [DOI: 10.1007/s10067-010-1616-y]
  56. Sheu SJ. Endophthalmitis. *Korean J Ophthalmol.* 2017; **31(4)**: 283-289. [PMID: 28752698]; [DOI: 10.3341/kjo.2017.0036]
  57. Callegan MC, Engelbert M, Parke DW, Jett BD, Gilmore MS. Bacterial endophthalmitis: epidemiology, therapeutics, and bacterium-host interactions. *Clin Microbiol Rev.* 2002; **15(1)**: 111-24. [PMID: 11781270]; [DOI: 10.1128/cmr.15.1.111-124.2002]
  58. Abu El-Asrar AM, Al-Amro SA, Al-Mosallam AA, Al-Obeidan S. Post-traumatic endophthalmitis: causative organisms and visual outcomes. *Eur J Ophthalmol.* 1999; **9**: 21-31. [PMID: 10230588]; [DOI: 10.1177/112067219900900104]
  59. Durand ML. Endophthalmitis. *Clin Microbiol Infect.* 2013; **19(3)**: 227-34. [PMID: 23438028]; [DOI: 10.1111/1469-0691.12118]
  60. Durand ML. Bacterial endophthalmitis. *Curr Infect Dis Rep.* 2009 Jul; **4**: 283-8. [PMID: 19545497]; [DOI: 10.1007/s11908-009-0042-2]

61. Harry F, Shah V. Outline of Postoperative Endophthalmitis. Review of Endophthalmitis Following Cataract Surgery: Prophylaxis and Treatment. Report, Bascom Palmer Eye Institute, August, 2015.
62. Lalwani GA, Flynn HW, Scott IU, Quinn CM, Berrocal AM, Davis JL, Murray TG, Smiddy WE, Miller D. Acute-onset endophthalmitis after clear corneal cataract surgery (1996-2005). Clinical features, causative organisms, and visual acuity outcomes. *Ophthalmol.* 2008; **115**(3): 473-6. [PMID: 18067969]; [DOI: 10.1016/j.ophtha.2007.06.006]
63. Jackson TL, Eykyn SJ, Graham EM, Stanford MR. Endogenous bacterial endophthalmitis: a 17-year prospective series and review of 267 reported cases. *Surv Ophthalmol.* 2003; **48**: 403-423. [PMID: 12850229]; [DOI: 10.1016/s0039-6257(03)00054-7]
64. Al-omran AM, Abboud EB, Abu el-asrar AM. Microbiologic spectrum and visual outcome of posttraumatic endophthalmitis. *Retina.* 2007; **27**(2): 236-42. [PMID: 17290207]; [DOI: 10.1097/01.iae.0000225072.68265.ee]
65. Ramadan RT, Ramirez R, Novosad BD, Callegan MC. Acute inflammation and loss of retinal architecture and function during experimental *Bacillus* endophthalmitis. *Curr Eye Res.* 2006; **31**: 955-965. [PMID: 17114121]; [DOI: 10.1080/02713680600976925]
66. Comer RM, Dawson E, Plant G, Acheson JF, Lee JP. Causes and outcomes for patients presenting with diplopia to an eye casualty department. *Eye.* 2007; **21**(3): 413-8. [PMID: 16732215]; [DOI: 10.1038/sj.eye.6702415]
67. Toro J, Burbano LE, Reyes S, Barreras P. Cavernous sinus syndrome: need for early diagnosis. *BMJ Case Rep.* 2015; **2015**. [PMID: 25819816]; [DOI: 10.1136/bcr-2014-206999]
68. Fu W, Duan L, Geng S. Surgical Outcomes of Cavernous Sinus Syndrome in Pituitary Adenomas. *World Neurosurg.* 2017; **107**: 526-533. [PMID: 28826864]; [DOI: 10.1016/j.wneu.2017.08.049]
69. van Gijn J, Kerr RS, Rinkel GJ. Subarachnoid haemorrhage. *Lancet.* 2007 Jan; **369**(9558): 306-18. [PMID: 17258671]; [DOI: 10.1016/S0140-6736(07)60153-6]
70. Nieuwkamp DJ, Setz LE, Algra A, Linn FHH, de Rooij NK, Rinkel GJ. Changes in case fatality of aneurysmal subarachnoid haemorrhage over time, according to age, sex, and region: a meta-analysis. *Lancet Neurol.* 2009; **8**(7): 635-42. [PMID: 19501022]; [DOI: 10.1016/S1474-4422(09)70126-7]
71. Wilne SH, Ferris RC, Nathwani A, Kennedy CR. The presenting features of brain tumours: a review of 200 cases. *Arch Dis Child.* 2006; **91**(6): 502-6. [PMID: 16547083]; [DOI: 10.1136/adc.2005.090266]
72. Naggara ON, Lecler A, Oppenheim C, Meder JF, Raymond J. Endovascular treatment of intracranial unruptured aneurysms: a systematic review of the literature on safety with emphasis on subgroup analyses. *Radiol.* 2012; **263**: 828-835. [PMID: 22623696]; [DOI: 10.1148/radiol.12112114]
73. Wilne S, Collier J, Kennedy C, Jenkins A, Grout J, Mackie S, Koller K, Grundy R, Walker D. Progression from first symptom to diagnosis in childhood brain tumours. *Eur J Pediatr.* 2012 Jan. **171**(1): 87-93. [PMID: 21594769]; [DOI: 10.1007/s00431-011-1485-7]
74. Lanphear J, Sarnaik S. Presenting symptoms of pediatric brain tumors diagnosed in the emergency department. *Pediatr Emerg Care.* 2014 Feb; **2**: 77-80. [PMID: 24457493]; [DOI: 10.1097/PEC.0000000000000074]
75. Mehta V, Chapman A, Mceely PD, Walling S, Howes WJ. Latency between symptom onset and diagnosis of pediatric brain tumors: an Eastern Canadian geographic study. *Neurosurg.* 2002; **51**(2): 365-72. [PMID: 12182774]
76. Haimi M, Peretz nahum M, Ben arush MW. Delay in diagnosis of children with cancer: a retrospective study of 315 children. *Pediatr Hematol Oncol.* 2004; **21**(1): 37-48. [PMID: 14660305]
77. Halperin EC, Friedman HS. Is there a correlation between duration of presenting symptoms and stage of medulloblastoma at the time of diagnosis?. *Cancer.* 1996; **78**(4): 874-80. [PMID: 8756384]; [DOI: 10.1002/(SICI)1097-0142(19960815)78:4<874::AID-CNCR26>3.0.CO;2-R]
78. Vaziri K, Schwartz SG, Flynn HW, Kishor KS, Moshfegi AA. Eye-related Emergency Department Visits in the United States, 2010. *Ophthalmol.* 2016; **123**(4): 917-9. [PMID: 26704884]; [DOI: 10.1016/j.ophtha.2015.10.032]