

The Age Significance of Lumbar Disc Extrusion or Sequestration: A Systematic Review of the Literature

Sombat Kunakornsawat, Thavat Prasarithra, Pritsanai Pruttikul, Chaiwat Piyaskulkaew, Tinnakorn Pluemvitayaporn

Sombat Kunakornsawat, Thavat Prasarithra, Pritsanai Pruttikul, Chaiwat Piyaskulkaew, Tinnakorn Pluemvitayaporn, Center of excellence in orthopedic, Lerdsin General Hospital, Bangkok, Thailand

Correspondence to: Sombat Kunakornsawat, Center of excellence in orthopedic, 9th floor, Lerdsin General Hospital, 190 Silom Rd. Bangrak; Bangkok, 10500, Thailand

Email: batkunakornsawat@gmail.com

Telephone: +66-81-6457671

Received: April 16, 2015

Revised: June 6, 2015

Accepted: June 9, 2015

Published online: October 23, 2015

© 2015 The Authors. Published by ACT Publishing Group Ltd.

Key words: Age; Lumbar disc; Extrusion; Sequestration

Kunakornsawat S, Prasarithra T, Pruttikul P, Piyaskulkaew C, Pluemvitayaporn T. The Age Significance of Lumbar Disc Extrusion or Sequestration: A Systematic Review of the Literature. *International Journal of Orthopaedics* 2015; 2(5): 421-426 Available from: URL: <http://www.ghrnet.org/index.php/ijo/article/view/1261>

ABSTRACT

BACKGROUND CONTEXT: Surgery of lumbar disc extrusion or sequestration is usually reported in young adult. The significance of age has not been determined when one considers that lumbar disc herniation is the most common reason of spine surgery.

AIM: To review the recent evidence and document the mean age at the time of lumbar disc surgery.

METHOD: A search of Medline, Embase, Cochrane database and cross references from 1980 to 2014 was conducted to identify the mean age of patients who underwent lumbar disc surgery. Two reviewers independently evaluated and extracted English published articles that contained keywords of lumbar disc herniation, extrusion and sequestration. Quality of evidence was classified into 3 levels.

RESULTS: There were totally 134 papers, 49 were excluded. Of all the 11,803 patients, 7,021 were male (59.48%) and 4,782 were female (40.52%) respectively with mean age of 41.75 years, age ranged from 18 to 96 years. All 85 articles were classified as 19 randomized controlled none concealed trials - level I, 34 prospective comparative studies - level II and 32 retrospective - level III evidence.

CONCLUSION: Patients with proven damaged disc(s) are at greater risk to have lumbar discectomy at the age of 41.75 years.

INTRODUCTION

The human lumbar intervertebral disc is a complex structure with three distinct components that undergoes progressive changes with age, chemical and mechanical process^[1-4]. In lumbar disc literature, the advanced damage of extrusion or sequestration may inevitably result in other spinal deformities and potential serious consequences^[5-7]. Tears within the disc can present itself at any age with some exceptional cases in children^[8-9]. Even since the first description of the ruptured intervertebral disc in 1934^[10], the presence of well established tissue damage has been widely used as an indication requiring surgical treatment. There are now an increasing number of scientific articles using modern technology such as intervertebral disc tissue engineering, advance biochemistry assay method and cell therapy^[11-16]. Surprisingly, the influence of range or mean age when patients met the criteria for surgery has not been raised. For the purposes of this study, the review was conducted to document the mean age at the time of surgery as well as its significance and emphasize some interesting information in the literature.

MATERIALS AND METHODS

Relevant randomized controlled trials (RCTs) and comparative observational studies were identified using computerized and hard

document search. Most common database of published literature included the Cochrane database, Embase and Medicine through Pub Med and cross references. Search terms or keywords were lumbar disc herniation, extrusion and sequestration. Only English language articles between 1980 and 2014 that addressed adult patients older than 18 years, range and the mean age at the time of surgery, at a single level and no other spine conditions were included. Studies not involving surgery such as non-operative treatment or laboratory, radiology, case report and duplicated reports were discarded.

Data from the included studies were extracted by 2 independent reviewers using a standard data abstraction sheet. The relevant information were (1) Country / department; (2) study design (prospective or retrospective); (3) types of surgery: a. standard with or without microscope and b. minimal invasive surgery, MIS (percutaneous or endoscopic discectomy); (4) number and gender of patients and (5) mean and range of age. Selection was based on the basis of title, abstract and sufficient detail in the materials and methods. Disagreement was resolved by discussion and clarified information.

Quality of evidence was graded into 3 levels^[17]: Level I - prospective randomized controlled trial (RCT); Level II - prospective non-randomized comparative study; Level III - retrospective comparative studies or case-controlled study. Level IV - case series or reports and Level V - expert opinions were excluded. For quality of RCT, methodology should contain detail of randomization, concealment and clear description of group allocation^[18].

RESULTS

There were totally 134 papers (Table 1), 49 were excluded as they did not include relevant data. One quarter (25%) of these contained incomplete data of age and gender. Of the 85 accepted reports, there were 19 randomized controlled none concealed trials - level I^[19-37], 34 prospective comparative studies - level II^[38-71] and 32 retrospective - level III evidence^[72-103]. Of these, 35 articles were from the United States, 23 from Europe and 23 from Asia while 4 came from other countries.

Among physicians, orthopedists (45 articles) were more frequent than neurosurgeons (33 articles). Co-ordination between orthopedist and neurosurgeon was found in 7 papers. There were totally 11,803 patients, 7,021 male (59.48%) and 4,782 female (40.52%) with mean age of 41.75 years, range 18-96 years.

Patients were operated on using standard open discectomy (56 articles) - with microscope 30 and without 40. Among those with minimal invasive surgery, there were 34 endoscopic, 12 percutaneous and 29 studies that used more than 1 technique for comparison.

DISCUSSION

Available literature has almost dealt with patient benefit for surgery and which surgical technique that would provide the best outcomes.

Range of year	Papers
1981-1985	9
1986-1990	13
1991-1995	14
1996-2000	13
2001-2005	25
2006-2010	38
2011-2014	21
Total articles published at 5 years intervals.	134

The review at present shows the strong increase in overall rate of lumbar disc surgery. Findings indicate high expected growing numbers of first disc surgery with greater rate found in the United States than other continents^[104,105]. Findings also reviewed a large number of patients - 11,803 patients underwent a variety of surgery performed by both Orthopaedists and Neurosurgeons at a range of academic and other spinal centers. The less frequent number of operations performed by neurosurgeons is not consistent with some earlier studies^[106-109].

Particularly in the last decade (twentieth century), the review clearly shows the burgeoning of new published articles as well as the emergence of less traumatic operative techniques in disc surgery^[110-111]. Despite an immense scientific interest, there are small number of RCTs. As expected, these trials are more difficult to design and conduct though a reasonable level of evidence can be provided. Within all studies a number of flaws such as high cross over rates, varied statistic techniques and unreported refusal rate could be considered as threats to their validity^[110,112]. The greater number of observation studies in this analysis may provide some accurate mean effects of one treatment compared with another. As has been stated, the strength of these studies from multicenters may be readily obtained when possible bias can be reduced^[112].

As shown, all the surgical techniques with results were quite variable. These national and international discrepancies may be due to professional preferences in choice of procedure differences in patient selection, study protocols and the uncertainty of definition and criteria for disc surgery^[106,108]. Open discectomy remains the most common approach for lumbar discectomy. A more acceptable concept of minimal invasive surgery (MIS) could be observed without providing comprehensive information regarding the potential risks. Part of recent meta-analysis also concluded that better benefit could be obtained when performed in specialized centers by experienced surgeons^[113]. A consensus of opinion in the literature concerning the growing number of lumbar spine surgeries would highlight the need for further improvement that would maximize clinical outcomes.

For the significance of demographic data, the analysis found a greater frequency of lumbar disc surgery in men than in women without covering its effect on the prognosis. For the effect of age in disc literature, high frequency tears within the disc would appear to occur before the age 30 years and become more pronounced with advanced age^[114-116]. With regard to the mean age as present (41.75 years), the disc extrusion or sequestration is neither age dependent nor directly correlated with degeneration grade^[106-109]. At this period, some vulnerable patients may be subjected to a complex pattern of loads, causing an abrupt displacement of disc material that requires removal. Although the study could not substantiate the significance of age at the time of surgery, this mean value may be used as a considerable limit but not an objective criteria. As the displaced disc tissue can be resolved with time and certain risk factors, some patients who can tolerate their symptoms beyond this period may be treated without surgery.

To select a better option of treatment for a young patient with moderate nerve root compression is especially difficult. Surgeons should be realized firstly that despite a successful discectomy, some patients may have the probability of recurrent pain and disability. Secondly, there are countless patients who undergoing disc re-operation. Literature indicated that more than half may need re-operation within 0.5-1 year after primary surgery^[106-109]. Thirdly, some may have negative long-term results irrespective of the method used to provide treatment.

On the basis of the current results, some patients may turn to in

moment of crisis at their middle age of life^[117], particularly after failed surgery. Adverse event of disc lesion as well as surgical procedure would also have devastating impact on families, societies and economics.

For lumbar disc herniation, non-surgical treatment should be recommended in the first instance^[118,119]. Discectomy that may give good results should mainly be restricted to clearly defined disc damage causing intractable pain and associated nerve entrapment signs^[120,121]. Patients should be fully informed of the potential risks of the first disc surgery as well as possibility of a later necessity for re-operation. Surgery should be justified to those who need it.

CONCLUSION

A large number of patients (11,803) with a male predominance of 59.44%, underwent a variety of surgeries, at the mean age of 41.75 years. There is a strong increase in overall rate of lumbar disc surgery performed by both Orthopaedists and Neurosurgeons. Open discectomy was the most common approach for lumbar discectomy.

CONFLICT OF INTERESTS

There are no conflicts of interest with regard to the present study.

REFERENCES

- Andersson GB, An HS, Oegema TR Jr, et al. Intervertebral disc degeneration. Summary of an AAOS/NIH/ORS workshop, September 2005. *J Bone Joint Surg Am* 2006;88:895-9.
- Fardon DF, Milette PC; Combined Task Forces of the North American Spine Society, American Society of Spine Radiology, and American Society of Neuroradiology. Nomenclature and classification of lumbar disc pathology. Recommendations of the Combined task Forces of the North American Spine Society, American Society of Spine Radiology, and American Society of Neuroradiology. *Spine (Phila Pa 1976)*. 2001 Mar 1;26(5):E93-E113.
- Harada Y, Nakahara S. A pathologic study of lumbar disc herniation in the elderly. *Spine (Phila Pa 1976)* 1989;14:1020-4.
- Yasuma T, Koh S, Okamura T, et al. Histological changes in aging lumbar intervertebral discs. Their role in protrusions and prolapses. *J Bone Joint Surg Am* 1990;72:220-9.
- Saal JA. Natural history and nonoperative treatment of lumbar disc herniation. *Spine (Phila Pa 1976)* 1996;21:2S-9S.
- Boos N, Weissbach S, Rohrbach H, et al. Classification of age-related changes in lumbar intervertebral discs: 2002 Volvo Award in basic science. *Spine (Phila Pa 1976)* 2002; 27:2631-44.
- Vernon-Roberts B, Moore RJ, Fraser RD. The natural history of age-related disc degeneration: the pathology and sequelae of tears. *Spine (Phila Pa 1976)* 2007;32:2797-804.
- Martínez-Lage JF, Fernández Cornejo V, López F, et al. Lumbar disc herniation in early childhood: case report and literature review. *Childs Nerv Syst* 2003;19:258-60.
- Hicks GE, Morone N, Weiner DK. Degenerative lumbar disc and facet disease in older adults: prevalence and clinical correlates. *Spine (Phila Pa 1976)* 2009;34:1301-6.
- Mixter WJ, Barr JS. Rupture of the intervertebral disc with involvement of the spinal canal. *N Engl J Med* 1934;211:210-215.
- Takeno K, Kobayashi S, Negoro K, Uchida K, Miyazaki T, Yayama T, Shimada S, Baba H. Physical limitations to tissue engineering of intervertebral disc cells: effect of extracellular osmotic change on glycosaminoglycan production and cell metabolism. Laboratory investigation. *J Neurosurg Spine*. 2007 Dec;7(6):637-44.
- Masuda K. Biological repair of the degenerated intervertebral disc by the injection of growth factors. *Eur Spine J*. 2008 Dec;17 Suppl 4:441-51.
- Kandel R, Roberts S, Urban JP. Tissue engineering and the intervertebral disc: the challenges. *Eur Spine J*. 2008 Dec;17 Suppl 4:480-91.
- Sobajima S, Vadala G, Shimer A, Kim JS, Gilbertson LG, Kang JD. Feasibility of a stem cell therapy for intervertebral disc degeneration. *Spine J*. 2008 Nov-Dec;8(6):888-96. Epub 2007 Dec 21.
- Hohaus C, Ganey TM, Minkus Y, Meisel HJ. Cell transplantation in lumbar spine disc degeneration disease. *Eur Spine J*. 2008 Dec;17 Suppl 4:492-503.
- Rodriguez AG, Slichter CK, Acosta FL, et al. Human disc nucleus properties and vertebral endplate permeability. *Spine (Phila Pa 1976)* 2011;36:512-20.
- Wright JG, Swiontkowski MF, Heckman JD. Introducing levels of evidence to the journal. *J Bone Joint Surg Am* 2003;85:1-3.
- Van Tulder M, Furlan A, Bombardier C, Bouter L. Updated method guidelines for systematic reviews in the Cochrane collaboration back review *Spine (Phila Pa 1976)*. 2003 Jun 15;28(12):1290-9.
- Hermantin FU, Peters T, Quartararo L, Kambin P. A prospective, randomized study comparing the results of open discectomy with those of video-assisted arthroscopic microdiscectomy. *J Bone Joint Surg Am*. 1999 Jul;81(7):958-65.
- Pookarnjanamorakot C, Laohacharoensombat W, Jaovisidha S. The clinical efficacy of Piroxicam fast-dissolving dosage form for postoperative pain control after simple lumbar spine surgery: a double-blinded randomized study. *Spine (Phila Pa 1976)*. 2002 Mar 1;27(5):447-51.
- U. Schick, J. Döhnert, A. Richter, A. König, H. E. Vitzthum: Microendoscopic lumbar discectomy versus open surgery: an intraoperative EMG study. *Eur Spine J*. 2002; 11:20–26
- Ryuichi Sasaoka, Hiroaki Nakamura, Sadahiko Konishi, Ryuichi Nagayama, Eisuke Suzuki, Hidetomi Terai, Kunio Takaoka: Objective assessment of reduced invasiveness in MED Compared with conventional one-level laminotomy. *Eur Spine J*. 2006; 15: 577–582.
- Osterman H, Seitsalo S, Karppinen J, Malmivaara A. Effectiveness of microdiscectomy for lumbar disc herniation: a randomized controlled trial with 2 years of follow-up. *Spine (Phila Pa 1976)*. 2006 Oct 1;31(21):2409-14.
- Hoogland T, Schubert M, Miklitz B, Ramirez A. Transforaminal posterolateral endoscopic discectomy with or without the combination of a low-dose chymopapain: a prospective randomized study in 280 consecutive cases. *Spine (Phila Pa 1976)*. 2006 Nov 15;31(24):E890-7.
- Katayama Y, Matsuyama Y, Yoshihara H, Sakai Y, Nakamura H, Nakashima S, Ito Z, Ishiguro N. Comparison of surgical outcomes between macro discectomy and microdiscectomy for lumbar disc herniation: a prospective randomized study with surgery performed by the same spine surgeon. *J Spinal Disord Tech*. 2006 Jul;19(5):344-7.
- Righesso O, Falavigna A, Avanzi O. Comparison of open discectomy with microendoscopic discectomy in lumbar disc herniations: results of a randomized controlled trial. *Neurosurgery*. 2007 Sep;61(3):545-9.
- Ryang YM, Oertel MF, Mayfrank L, Gilsbach JM, Rohde V. Standard open microdiscectomy versus minimal access trocar microdiscectomy: results of a prospective randomized study. *Neurosurgery*. 2008 Jan;62(1):174-81;
- Ruetten S, Komp M, Merk H, Godolias G. Full-endoscopic interlaminar and transforaminal lumbar discectomy versus conventional microsurgical technique: a prospective, randomized, controlled study. *Spine (Phila Pa 1976)*. 2008 Apr 20;33(9):931-9.
- Marko Brock, Philip Kunkel, Luca Papavero Lumbar microdiscectomy: subperiosteal versus transmuscular approach and influence on the early postoperative analgesic consumption. *Eur Spine J* (2008) 17:518–522.

- 30 Martin Barth, Christel Weiss, Claudius Thome. Two-Year Outcome After Lumbar Microdiscectomy Versus Microscopic Sequestrectomy Part I: Evaluation of Clinical Outcome. *Spine (Phila Pa 1976)* 2008 Feb 1; 33(3): 265-72.
- 31 Arts MP, Brand R, van den Akker ME, Koes BW, Bartels RH, Peul WC; Leiden-The Hague Spine Intervention Prognostic Study Group (SIPS). Tubular discectomy vs conventional microdiscectomy for sciatica: a randomized controlled trial. *JAMA*. 2009 Jul 8;302(2):149-58.
- 32 Jorg Franke, R. Greiner-Perth, H. Boehm, K. Mahlfeld, H. Grasshoff, Y. Allam, F. Awiszus. Comparison of a minimally invasive procedure versus standard microscopic discectomy: a prospective randomized controlled clinical trial. *Eur Spine J* (2009) 18:992–1000.
- 33 Marco Teli, Alessio Lovi, Marco Brayda-Bruno, Antonino Zagra, Andrea Corriero, Fabrizio Giudici, Leone Minoia. Higher risk of dural tears and recurrent herniation with lumbar micro-endoscopic discectomy. *Eur Spine J* (2010) 19:443–450.
- 34 Bhavuk Garg, Upendra Bidre Nagraja, Arvind Jayaswal. Microendoscopic versus open discectomy for lumbar disc herniation: a prospective randomised study. *Journal of Orthopaedic Surgery* 2011;19(1):30-4.
- 35 Alexander Bailey, MD, Ali Araghi, DO, Scott Blumenthal, MD, George V. Huffmon, MD, and the Annular Repair Clinical Study Group. Prospective, Multicenter, Randomized, Controlled Study of Annular Repair in Lumbar Discectomy. *Spine Volume 38, Number 14, pp 1161–1169*.
- 36 Mohamed Hussein, Ashraf Abdeldayem, Mahmoud M. M. Mattar. Surgical technique and effectiveness of microendoscopic discectomy for large uncontained lumbar disc herniations: a prospective, randomized, controlled study with 8 years of follow-up. *Eur Spine J* (2014) 23:1992–1999.
- 37 Peul WC, van Houwelingen HC, van den Hout WB, Brand R, Eekhof JA, Tans JT, Thomeer RT, Koes BW. Surgery versus prolonged conservative treatment for sciatica. *N Engl J Med*. 2007 May 31;356(22):2245-56.
- 38 Dan M Spengler, MD: Lumbar Discectomy: Results With Limited Disc Excision and Selective Foraminotomy. *Spine*. 1982;7 (6): 604-607.
- 39 S B Tibrewal, M J Pearcy, I Portek, J Spivey: A Prosective Study of Lumbar Spinal Movements Before and After Discectomy Using Biplanar Radiography: Correlation of Clinical and Radiographic Findings. *Spine*. 1985; 10 (5): 455-460.
- 40 Roger P Jackson, James J Glah: Foraminal and Extraforaminal Lumbar Disc Herniation: Diagnosis and Treatment. *Spine*. 1987; 12 (6): 577-585.
- 41 Neil Kahanovitz, Kathleen Viola and John Muculloch: Limited Surgical Discectomy and Microdiscectomy: A clinical Comparison. *Spine*. 1989; 14 (1): 79-81.
- 42 Stern MB. Early experience with percutaneous lateral discectomy. *Clin OrthopRelat Res*. 1989 Jan;(238):50-5.
- 43 C Edmund Graham: Percutaneous Posterolateral Lumbar Discectomy. *Clin Orthop Relat Res*. 1989 Jan;(238):104-106.
- 44 Parviz Kambin, Jonathan L Schaffer: Percutaneous Lumbar Discectomy: Review of 100 Patients and Current Practice. *Clin Orthop Relat Res*. 1989 Jan;(238):24-34.
- 45 Neil Kahanovitz, Kathleen Viola, theodore Goldstein and Edgar Dawson: A Multicenter Analysis of Percutaneous Discectomy. *Spine*. 1990; 15 (7): 713-716.
- 46 Jonathan L Schaffer and Parviz Kambin: Percutaneous Posterolateral Lumbar Discectomy and Decompression with a 6.9-Millimeter Cannula. *The Journal of Bone and Joint Surgery*.1991;73-A (6): 822-831.
- 47 Shigeru Hirabayashi, Kiyoshi Kumano, Yutaka Ogawa, Yoichi Aota and Susumu Maehiro: Microdiscectomy and Second Operation for Lumbar Disc Herniation. *Spine*.1993;18 (15):2206-2211.
- 48 Takashi Sakou and Akitoshi Masuda: Percutaneous Discectomy for Lumbar Disc Herniation: A Preliminary Report. *Clin Orthop Relat Res*. 1993 Jan;(286): 174-179.
- 49 Gus J Slotman and Sherman C Stein: Laparoscopic Lumbar Discectomy: Preliminary Report of a Minimally Invasive Anterior Approach to the Herniated L5-S1 Disk. *Surgical Laparoscopy and Endoscopy*. 1995;5 (5):363-369.
- 50 Hallett H Mathews, Martin T Evans, Harry J Molligan and Brenda H Long: Laparoscopic Discectomy With Anterior Lumbar Interbody Fusion: A Preliminary Review. *Spine*. 1995; 20, (16):1797-1802.
- 51 Michael A MacKay, Jeffrey S Fischgrund, Harry N Herkowitz, Lawrence T Kurz, Brian Hecht and Michael Schwartz: The Effect of Interposition Membrane on the Outcome of Lumbar Laminectomy and Discectomy. *Spine*. 1995; 20 (16):1793-1796.
- 52 M P Grevitt, A McLaren, I M Shackleford, R C Mulholland: Automated Percutaneous lumbar discectomy. *The Journal of Bone and Joint Surgery (Br)*. 1995; 77-B:626-9.
- 53 Larry D Herron, Judith A Turner, Laura A Novell and Suzanne L Kreif: Patient Selection for Lumbar Discectomy With a Revised Objective Rating System. *Clin Orthop Relat Res*. 1996;(325): 148-155.
- 54 Türeyen K. One-level one-sided lumbar disc surgery with and without microscopic assistance: 1-year outcome in 114 consecutive patients. *J Neurosurg*. 2003 Oct;99(3 Suppl):247-50.
- 55 Fisher C, Noonan V, Bishop P, Boyd M, Fairholm D, Wing P, Dvorak M. Outcome evaluation of the operative management of lumbar disc herniation causing sciatica. *J Neurosurg (Spine 4)* 2004; 100:317–324.
- 56 P.C. Hsieh, C.H. Wang. Posterior endoscopic lumbar discectomy using a thoracoport as tubular retractor. *Minim Invas Neurosurg* 2004 Oct; 47(5): 319-23.
- 57 Kenneth M. Alò, Robert E. Wright, John Sutcliffe, Scott A. Brandt. Percutaneous Lumbar Discectomy: Clinical Response in an Initial Cohort of Fifty Consecutive Patients With Chronic Radicular Pain. *Pain Practice*, Vol. 4, Issue 1, 2004: 19–29.
- 58 D. D. Limbrick Jr., N. M. Wright. Verification of Nerve Root Decompression during Minimally–Invasive Lumbar Microdiscectomy: A Practical Application of Surgeon–Driven Evoked EMG. *Minim Invas Neurosurg* 2005 Oct; 48(5): 273-7.
- 59 Cagatay Ozturk, Mehmet Tezer, Mehmet Aydogan, Mercan Sarier, Azmi Hamzaoglu: Posterior endoscopic discectomy for the treatment of lumbar disc herniation. *Acta Orthop Belg.*, 2006, 72, 347-352.
- 60 Seungcheol Lee, Seok-Kang Kim, Sang-Ho Lee, Won Joong Kim, Won-Chul Choi, Gun Choi, Song-Woo Shin: Percutaneous endoscopic lumbar discectomy for migrated disc herniation: classification of disc migration and surgical approaches. *Eur Spine J* (2007) 16:431–437.
- 61 Kadir Kotil, Mustafa Akcetin, Turgay Bilge. A Minimally Invasive Transmuscular Approach to Far-lateral L5-S1 Level Disc Herniations A Prospective Study. *J Spinal Disord Tech* 2007 Apr; 20(2): 132-8.
- 62 Erich Kast, Joachim Oberle, Hans-Peter Richter, Wolfgang Borm. Success of Simple Sequestrectomy in Lumbar Spine Surgery Depends on the Competence of the Fibrous Ring A Prospective Controlled Study of 168 Patients. *Spine (Phila Pa 1976)* 2008 Jun 15; 33(14): 1567-71.
- 63 F Prochet, V Bartanusz, F S Kleinstueck, F Lattig, D Jeszenszky, D Grob, A F Mannion: Microdiscectomy compared with standard discectomy: an old problem revisited with new outcome measures within the framework of a spine surgical registry. *Eur Spine J* (2009) 18: 360-366.
- 64 Kotryna Veresciagina, Bronius Spakauskas, Kazys Vytautas Ambrozaitis. Clinical outcomes of patients with lumbar disc herniation, selected for one-level open-discectomy and microdiscectomy. *Eur Spine J* (2010) 19:1450–1458.
- 65 Akagi R, Aoki Y, Ikeda Y, Nakajima F, Ohtori S, Takahashi K,

- Yamagata M. Comparison of early and late surgical intervention for lumbar disc herniation: is earlier better? *J Orthop Sci.* 2010 May;15(3):294-8.
- 66 Yong Ahn, Sang-Ho Lee. Outcome predictors of percutaneous endoscopic lumbar discectomy and thermal annuloplasty for discogenic low back pain. *Acta Neurochir* (2010) 152:1695–1702.
- 67 Katarina Silverplats, B. Lind, B. Zoc̄ga, K. Halldin, M. Gellerstedt, H. Brisby, L. Rutberg. Clinical factors of importance for outcome after lumbar disc herniation surgery: long-term follow-up. *Eur Spine J* (2010) 19:1459–1467.
- 68 Kyung-Chul Choi, MD, Jin-Sung Kim, MD, Byung-Uk Kang, MD, Choon Dae Lee, MD, and Sang-Ho Lee, MD, PhD. Changes in Back Pain After Percutaneous Endoscopic Lumbar Discectomy and Annuloplasty for Lumbar Disc Herniation: A Prospective Study. *Pain Medicine* 2011; 12: 1615–1621.
- 69 Hsien-Te Chen, Chun-Hao Tsai, Shao-Ching Chao, Ting-Hsien Kao, Yen-Jen Chen, Horng-Chaung Hsu, Chiung-Chyi Shen, Hsi-Kai Tsou. Endoscopic discectomy of L5–S1 disc herniation via an interlaminar approach: Prospective controlled study under local and general anesthesia. *Surg Neurol Int* 2011;2:93.
- 70 Kamrul Ahsan, Najmus-Sakeb, Alamgir Hossain, Shahidul Islam Khan, MA Awwal: Discectomy for primary and recurrent prolapse of lumbar intervertebral discs. *Journal of Orthopaedic Surgery* 2012;20(1):7-10.
- 71 Atman Desai, MD, Kimon Bekelis, MD, Perry A. Ball, MD, Jon Lurie, MD, Sohail K. Mirza, MD, Tor D. Tosteson, DSc, Wenyan Zhao, MS James N. Weinstein, DO, MS. Spine Patient Outcomes Research Trial: Do Outcomes Vary Across Centers for Surgery for Lumbar Disc Herniation? *Neurosurgery* 2012; 71:833–843.
- 72 Larry D Herron, Homer C Pheasant: Bilateral Laminotomy and Discectomy for Segmental Lumbar Disc Disease: Decompression with Stability. *Spine.* 1983; 8 (1): 86-98.
- 73 William A Friedman: Percutaneous Discectomy: An Alternative to Chemonucleolysis. *Neurosurgery.* 1983; 13 (5):542-547.
- 74 Joseph C Maroon, Adnan Abla: Microdiscectomy versus Chemonucleolysis. *Neurosurgery.* 1985; 16 (5):644-649.
- 75 Larry D Herron, Judith Turner: Patient Selection for Lumbar Laminectomy and Discectomy with a Revised Objective Rating System. *Clin Orthop Relat Res.* 1985;(199): 145-152.
- 76 Weinstein J, Spratt KF, Lehmann T, McNeill T, Hejna W. Lumbar disc herniation. A comparison of the results of chemonucleolysis and open discectomy after ten years. *J Bone Joint Surg Am.* 1986 Jan;68(1):43-54.
- 77 Zahrawi F. Microlumbar discectomy (MLD). *Spine (Phila Pa 1976).* 1988 Mar;13(3):358-9.
- 78 Ramirez LF, Thisted R. Complications and demographic characteristics of patients undergoing lumbar discectomy in community hospitals. *Neurosurgery.* 1989 Aug;25(2):226-30.
- 79 Kahanovitz N, Viola K, Gallagher M. Long-term strength assessment of postoperative discectomy patients. *Spine (Phila Pa 1976).* 1989 Apr;14(4):402-3.
- 80 Caspar W, Campbell B, Barbier DD, Kretschmmer R, Gotfried Y. The Caspar microsurgical discectomy and comparison with a conventional standard lumbar disc procedure. *Neurosurgery.* 1991 Jan;28(1):78-86.
- 81 Matsunaga S, Sakou T, Taketomi E, Ijiri K. Comparison of operative results of lumbar disc herniation in manual laborers and athletes. *Spine (Phila Pa 1976).* 1993 Nov;18(15):2222-6.
- 82 Kowalski JM, Olsewski JM, Simmons ED Jr. Results of intervertebral discectomy without fusion at L4-5 versus L5-S1. *J Spinal Disord.* 1995 Dec;8(6):457-63.
- 83 Knop-Jergas BM, Zucherman JF, Hsu KY, DeLong B. Anatomic position of a herniated nucleus pulposus predicts the outcome of lumbar discectomy. *J Spinal Disord.* 1996 Jun;9(3):246-50.
- 84 Stambough JL. Lumbar disk herniation: an analysis of 175 surgically treated cases. *J Spinal Disord.* 1997 Dec;10(6):488-92.
- 85 Ditsworth DA. Endoscopic transforaminal lumbar discectomy and reconfiguration: a postero-lateral approach into the spinal canal. *Surg Neurol.* 1998 Jun;49(6):588-97.
- 86 Yorimitsu E, Chiba K, Toyama Y, Hirabayashi K. Long-term outcomes of standard discectomy for lumbar disc herniation: a follow-up study of more than 10 years. *Spine (Phila Pa 1976).* 2001 Mar 15;26(6):652-7.
- 87 Wenger M, Mariani L, Kalbarczyk A, Gröger U. Long-term outcome of 104 patients after lumbar sequestrectomy according to Williams. *Neurosurgery.* 2001 Aug;49(2):329-34.
- 88 Tsou PM, Yeung AT. Transforaminal endoscopic decompression for radiculopathy secondary to intracanal noncontained lumbar disc herniations: outcome and technique. *Spine J.* 2002 Jan-Feb;2(1):41-8.
- 89 Nakagawa H, Kamimura M, Uchiyama S, Takahara K, Itsubo T, Miyasaka T. Microendoscopic discectomy (MED) for lumbar disc prolapse. *J Clin Neurosci.* 2003 Mar;10(2):231-5.
- 90 Morgan-Hough CV, Jones PW, Eisenstein SM. Primary and revision lumbar discectomy. A 16-year review from one centre. *J Bone Joint Surg Br.* 2003 Aug;85(6):871-4.
- 91 Tzaan WC. Transforaminal percutaneous endoscopic lumbar discectomy. *Chang Gung Med J.* 2007 May-Jun;30(3):226-34.
- 92 Mariconda M, Galasso O, Secondulfo V, Rotonda GD, Milano C. Minimum 25-year outcome and functional assessment of lumbar discectomy. *Spine (Phila Pa 1976).* 2006 Oct 15;31(22):2593-9.
- 93 Jang JS, An SH, Lee SH. Transforaminal percutaneous endoscopic discectomy in the treatment of foraminal and extraforaminal lumbar disc herniations. *J Spinal Disord Tech.* 2006 Jul;19(5):338-43.
- 94 Wu X, Zhuang S, Mao Z, Chen H. Microendoscopic Discectomy for Lumbar Disc Herniation Surgical Technique and Outcome in 873 Consecutive Cases. *Spine.* 2006;31(23):2689-2694.
- 95 J. F. Harrington, P. French. Open versus Minimally Invasive Lumbar Microdiscectomy: Comparison of Operative Times, Length of Hospital Stay, Narcotic Use and Complications. *Minim Invas Neurosurg* 2008 Feb;51(1):30-5.
- 96 Kyeong-seong Yeom, Yong-soo Choi. Full endoscopic contralateral transforaminal discectomy for distally migrated lumbar disc herniation. *J Orthop Sci* (2011) 16:263–269.
- 97 Lau D, Han SJ, Lee JG, Lu DC, Chou D. Minimally invasive compared to open micro discectomy for lumbar disc herniation. *J Clin Neurosci* 2011 Jan; 18(1): 81-4.
- 98 Kyung-Chul Choi, Jin Sung Kim, Kyeong Sik Ryu, Byung Uk Kang, Yong Ahn, Sang-Ho Lee: Percutaneous Endoscopic Lumbar Discectomy for L5-S1 Disc Herniation: Transforaminal vs Interlaminar approach. *Pain Physician* 2013; 16:547-556.
- 99 Hongwei Wang & Bo Huang & Wenjie Zheng & Changqing Li & Zhengfeng Zhang & Jian Wang & Yue Zhou. Comparison of early and late percutaneous endoscopic lumbar discectomy for lumbar disc herniation. *Acta Neurochir* (2013) 155:1931–1936.
- 100 Haidong Xu, Xiaozhou Liu, Gang Liu, Jiangning Zhao, Qiang Fu, Bin Xu. Learning Curve of Full-Endoscopic Technique Through Interlaminar Approach for L5/S1 Disk Herniations. *Cell Biochem Biophys* (2014) 70:1069–1074.
- 101 Mitsunori Yoshimoto, MD, Takehito Iwase, MD, Tsuneo Takebayashi, MD, Kazunori Ida, MD, and Toshihiko Yamashita, MD. Microendoscopic Discectomy for Far Lateral Lumbar Disk Herniation: Less Surgical Invasiveness and Minimum 2-Year Follow-Up Results. *J Spinal Disord Tech* 2014;27:E1–E7.
- 102 Alexander Aichmair, Jerry Y. Du, Jennifer Shue, Gisberto Evangelisti, Andrew A. Sama, Alexander P. Hughes, Darren R. Lebl, Jayme C. Burket, Frank P. Cammisa, Federico P. Girardi. Microdiscectomy for the Treatment of Lumbar Disc Herniation: An Evaluation of Reoperations and Long-Term Outcomes. *Evid Based Spine Care J* 2014;5:77–86.
- 103 Corriero OV, Morichi Rm Vagaggini A, Paoli L, Guizzardi G: Lumbar Herniated Disc Treated by Microdiscectomy Aline or Microdiscectomy Plus an Interlaminar Shock Absorbing Device: Retrospective Study with Minimum 3 Years Follow-Up. *Journal*

- of Spine: Advanced Techniques in Spine Surgery 2014; 3 (2).
- 104 Desai A, Bekelis K, Ball PA, Lurie J, Mirza SK, Tosteson TD, Zhao W, Weinstein JN. Spine patient outcomes research trial: do outcomes vary across centers for surgery for lumbar disc herniation? *Neurosurgery*. 2012 Oct;71(4):833-42.
 - 105 Yoshihara H, Yoneoka D. National trends in the surgical treatment for lumbar degenerative disc disease: United States, 2000 to 2009. *Spine J*. 2015 Feb 1;15(2):265-71.
 - 106 Keskimäki I, Seitsalo S, Osterman H, Rissanen P. Reoperations after lumbar disc surgery: a population-based study of regional and interspecialty variations. *Spine (Phila Pa 1976)*. 2000 Jun 15;25(12):1500-8.
 - 107 Cheng J, Wang H, Zheng W, Li C, Wang J, Zhang Z, Huang B, Zhou Y. Reoperation after lumbar disc surgery in two hundred and seven patients. *Int Orthop*. 2013 Aug;37(8):1511-7.
 - 108 Martin BI, Mirza SK, Flum DR, Wickizer TM, Heagerty PJ, Lenkoski AF, Deyo RA. Repeat surgery after lumbar decompression for herniated disc: the quality implications of hospital and surgeon variation. *Spine J*. 2012 Feb;12(2):89-97.
 - 109 Kim CH, Chung CK, Park CS, Choi B, Kim MJ, Park BJ. Reoperation rate after surgery for lumbar herniated intervertebral disc disease: nationwide cohort study. *Spine (Phila Pa 1976)*. 2013 Apr 1;38(7):581-90.
 - 110 Smith N, Masters J, Jensen C, Khan A, Sprowson A. Systematic review of microendoscopic discectomy for lumbar disc herniation. *Eur Spine J*. 2013 Nov;22(11):2458-65.
 - 111 Kamper SJ, Ostelo RW, Rubinstein SM, Nellensteijn JM, Peul WC, Arts MP, van Tulder MW. Minimally invasive surgery for lumbar disc herniation: a systematic review and meta-analysis. *Eur Spine J*. 2014 May;23(5):1021-43.
 - 112 Anderson PA, McCormick PC, Angevine PD. Randomized controlled trials of the treatment of lumbar disk herniation: 1983-2007. *J Am Acad Orthop Surg* 2008;16:566-7.
 - 113 Sidhu GS, Henkelman E, Vaccaro AR, Albert TJ, Hilibrand A, Anderson DG, Rihn JA. Minimally invasive versus open posterior lumbar interbody fusion: a systematic review. *Clin Orthop Relat Res*. 2014 Jun;472(6):1792-9.
 - 114 Kelsey JL, Githens PB, O'Conner T, et al. Acute prolapsed lumbar intervertebral disc. An epidemiologic study with special reference to driving automobiles and cigarette smoking. *Spine (Phila Pa 1976)* 1984;9:608-13.
 - 115 Mundt DJ, Kelsey JL, Golden AL, et al. An epidemiologic study of non-occupational lifting as a risk factor for herniated lumbar intervertebral disc. The Northeast Collaborative Group on Low Back Pain. *Spine (Phila Pa 1976)* 1993;18:595-602.
 - 116 Wolter T, Szabo E, Becker R, et al. Chronic low back pain: course of disease from the patient's perspective. *Int Orthop* 2011;35:717-24.
 - 117 Seyedmehdi M, Attarchi M, Ghaffari M, et al. Prognostic Factors for Return to Work After Low-Back Disc Herniation Surgery. *Asia Pac J Public Health* 2013;Jan 4.
 - 118 Ahn SH, Ahn MW, Byun WM. Effect of the transligamentous extension of lumbar disc herniations on their regression and the clinical outcome of sciatica. *Spine (Phila Pa 1976)* 2000;25:475-80.
 - 119 Sabuncuoglu H, Ozdogan S, Timurkaynak E. Spontaneous regression of extruded lumbar disc herniation: report of two illustrative case and review of the literature. *Turk Neurosurg* 2008;18:392-6.
 - 120 Carragee EJ, Han MY, Suen PW, et al. Clinical outcomes after lumbar discectomy for sciatica: the effects of fragment type and anular competence. *J Bone Joint Surg Am* 2003;85:102-8.
 - 121 Rajasekaran S, Bajaj N, Tubaki V, et al. ISSLS Prize winner: The anatomy of failure in lumbar disc herniation: an in vivo, multimodal, prospective study of 181 subjects. *Spine (Phila Pa 1976)* 2013;38:1491-500.
- Peer reviewer:** Santino Ottavio Tomasi, MD, Department of Neurosurgery, Christian-Doppler-Medical Center, Laboratory for Microsurgical Neuroanatomy, Paracelsus Medical University, Ignaz-Harrer-Str. 79, 5020 Salzburg, Austria.