

Outcome of Congenital Pseudoarthrosis Tibia Treated with Illizarov Fixator in Children

Muhammad Zubair Javaid¹, Junaid Khan², Muhammad Mohsin Javaid³, Zeeshan Zafar⁴, Imtiaz Ahmed Shakir⁵, Shahid Bashir⁶

1 Associate Professor Orthopaedics, Rawalpindi Medical University, Rawalpindi, Pakistan;

2 Registrar Orthopaedics, Rawalpindi Medical University, Rawalpindi, Pakistan;

3 Post-graduate resident Orthopaedics, Rawalpindi Medical University, Rawalpindi, Pakistan;

4 House Officer Orthopaedics, Rawalpindi Medical University, Rawalpindi, Pakistan;

5 Senior Registrar Orthopaedics, Rawalpindi Medical University, Rawalpindi, Pakistan;

6 Post-graduate resident Orthopaedics, Rawalpindi Medical University, Rawalpindi, Pakistan.

Conflict-of-interest statement: The author(s) declare(s) that there is no conflict of interest regarding the publication of this paper.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Correspondence to: Muhammad Zubair Javaid, Associate Professor Orthopaedics, Rawalpindi Medical University, Rawalpindi, Pakistan.

Email: drzubairortho@hotmail.com

Telephone: +92-333-5106566

Received: February 3, 2020

Revised: March 2, 2020

Accepted: March 10 2020

Published online: June 28, 2020

ABSTRACT

OBJECTIVES: To determine the outcome of congenital pseudoarthrosis tibia (CPT) managed with illizarov fixator.

METHODS: This prospective cross-sectional study was done from

20th March 2008 to 19th March 2018. All patients presenting to the Orthopaedic out patient department (OPD) with pseudoarthrosis tibia belonging to either gender aged up to 12 years were included. All patients underwent excision of tapering ends and illizarov fixator application. Outcome was assessed in terms of complications. Data regarding age, gender, side, type of CPT and any complication was noted and analyzed using SPSS version 23.

RESULTS: 16 patients were included in the study. 10 (62.5%) were males and 6 (37.5%) females. Mean age was 4 years. 10 (62.5%) patients were type-1 and 6 (37.5%) were type-2. Evidence of neurofibromatosis was seen in 9 (56.2%) patients. Most common complications were residual leg length discrepancy (LLD), mal-alignment and delayed consolidation which was seen in 8 (50%) patients.

CONCLUSION: CPT is a periosteal disease with variable results. The study concludes that good results are achieved in 50% patients.

Key Words: Congenital pseudoarthrosis tibia; Illizarov fixator; Leg length discrepancy; Neurofibromatosis

© 2020 The Author(s). Published by ACT Publishing Group Ltd. All rights reserved.

Javaid MZ, Khan J, Javaid MM, Zafar Z, Shakir IA, Bashir S. Outcome of Congenital Pseudoarthrosis Tibia Treated with Illizarov Fixator in Children. *International Journal of Orthopaedics* 2020; 7(3): 1289-1292 Available from: URL: <http://www.ghrnet.org/index.php/ijo/article/view/2792>

INTRODUCTION

Congenital pseudoarthrosis tibia (CPT) that may be obvious at birth or incipient, most commonly involves the distal half of tibia and often fibula in the same limb. Incidence is 1 in 250,000 live births^[1]. CPT is closely associated with neurofibromatosis that may not be the cause of pseudoarthrosis. Variable forms of fibrous dysplasia are also reported with CPT^[2]. True cause of poor healing at pseudoarthrosis is unknown, however few of the cases mentioned in literature are hamartomatous thickening of fibrous tissue, limited vascular potential, thickening of adherent periosteum and osteolytic fibromatosis^[3].

Natural history of CPT is neither predictable nor alterable with surgical and medical treatment. The treatment aims are to achieve

bony union at pseudoarthrosis, restore leg length discrepancy (LLD), prevent mal-alignment and to address ankle and foot problems^[4]. Different methods have been used to achieve the goal like intramedullary fixation, free fibular graft and illizarov technique and at times a combination of the above surgeries^[5]. Type of surgery to achieve union and prevent refracture remains controversial. Initial management is resection of entire pseudoarthrosis and surrounding hamartomatous tissue^[6]. Atrophic type of CPT is more challenging. Excision of the fibrous hamartoma, removing sclerotic tapering bone ends to sufficient cross sectional tubular bone, axial mal-alignment and ankle stabilization. Osseous union is difficult to achieve having a success rate of 31 to 56 percent^[7]. Union obtained may be temporary with 50% incidence of refracture^[8]. Secondary surgery for knee, ankle, foot, mal-alignment and LLD increases the frustration of surgeon and agony of patient to end up with amputation of limb^[9]. Involved limb is already short, removal of pathological tissue and tapering ends of bone further increases the limb-length discrepancy^[10].

Illizarov fixator is used for bone lengthening, compression at fracture site after generous excision of pathological bone and soft tissue^[1]. Valgus foot and ankle deformity is addressed with the same apparatus. Illizarov became available to us in early 1990's. Author experience with apparatus was boosted by 2006 earthquake. Illizarov can address pseudoarthrosis, LLD and foot component problems.¹ This technique is also useful when other modalities are exhausted. As there is scarcity of literature on CPT management and outcome, so this study was undertaken to determine the outcome of congenital pseudoarthrosis tibia (CPT) managed with illizarov fixator.

MATERIAL AND METHODS

This prospective cross-sectional study was done from 20th March 2008 to 19th March 2018 in the Department of Orthopaedics Benazir Bhutto Hospital and Holy Family Hospital, Rawalpindi, Pakistan. All patients presenting to the Orthopaedic out patient department (OPD) with pseudoarthrosis tibia belonging to either gender aged up to 18 years were included. Pre-existing infected CPT patients were excluded. Patients were assessed between 2008 to 2018 in Benazir Bhutto Hospital and Holy Family Hospital, which are attached hospitals of Rawalpindi Medical University. CPT was classified based on Boyd classification. All patients underwent excision of tapering ends and illizarov fixator application by the same consultant orthopaedic surgeon.

Illizarov procedure: Patients below the age of 5, only 3 rings were accommodatable; we used 1.5mm K-wire in these patients. Specially made sling for foot support was used made by the orthotic and prosthetic department, Rawalpindi Medical University, Rawalpindi. Hamartomatous tissue and tapering ends of bone were excised, till

Table 1 Complications.

S.No.	Complication	No. of patients	Percentage
1	Pin track infection	5	31.25%
2	Loosening, breakage and refracture of pin insertion site	4	25%
3	Neurovascular injury	1	6.25%
4	Temporary kinking of vessel	1	6.25%
5	Valgus deformity	3	18.75%
6	Residual LLD	8	50%
7	Refracture	4	25%
8	Non-compliance with bracing	10	62.50%
9	Diaphyseal malalignment	8	50%
10	Delayed consolidation	8	50%

good bleeding bones at least 70% of the shaft diameter was gained to increase contact surface area. Attempt was made to do compression at the fracture site, to avoid kinking of the vessel, a gap more than 4cm between ends per operation compression was avoided. Later this gap was reduced to 3cm as one patient of 2 years age got kinking of the vessel; which relieved after reopening the gap. Osteotomy of the proximal tibia with predrilling of osteotome was used. Distraction in younger children started at the 8th day and in older, 10th to 12th day. 1mm/day distraction in 0.25mm/6 hourly. For large bone defect, speed was reduced to 0.75mm/day. We kept the frame till good evidence of union at distraction and non-union site. For accurate approximation of middle and distal fragment, we used wire from heel to the middle fragment to rail road the middle fragment. Data was recorded on pre-formed proformas. Data analysis done using SPSS version 23.

RESULTS

A total of 16 patients were included in this study, 10 (62.5%) were males and 6 (37.5%) females. Age ranged from 2 to 17 years, with a mean age of 4 years. According to Boyd classification, 10 (62.5%) patients were type 1 and 6 (37.5%) were type 2. Evidence of neurofibromatosis was seen in 9 (56.2%) patients. LLD before surgery was seen in all patients and varied from 2 to 10 cm. 10 (62.5%) patients underwent primary surgery while 6 (37.5%) patients had already undergone more than two surgeries.

Most common complications were residual LLD, mal-alignment and delayed consolidation which was seen in 8 (50%) patients (Table 1).

Bone grafting was required in 3 (18.75%) patients. 1 patient aged 7 years had gained 1.5cm more limb length than the normal limb (Figure 1).

1 patient had unexplained resorption of the proximal bone at the site of osteotomy and distraction resulting in proximal pseudoarthrosis as well (Figure 2).

Another patient had a grossly deformed tibia at the age of 17 years. This patient was operated in early childhood with free non-vascularized fibular graft (Figure 3).

DISCUSSION

No standard surgical method for CPT has been defined. Basic treatment goal of these patients is to achieve union, prevent refracture, to correct LLD and correction of deformities particularly ankle joint^[1]. Different modes of treatment are clinically practiced like intramedullary fixation illizarov alone, illizarov with intramedullary fixation and free vascularized fibular graft^[12]. Age at which CPT should have surgery is quite controversial^[13]. Joseph et al believes surgery is beneficial at the age of 3 to 4 years while others has recommended the age of 5 to 6 years as having better bone stock. Children below the age of 5, bones are not only difficult to unite as low cross section area of the bone and most of them have early fractures^[14].

Illizarov technique has achieved union as well as addresses LLD, angular deformities and ankle deformities. This procedure has gained acceptance in many centers. Grill et al considered this procedure as gold standard for the treatment of CPT^[15]. Cho et al reported 20/23 patients with refracture. Late apical mal-alignment may develop even after union^[16]. One case of 11 year old female apparently normal looking proximal tibial bone after osteotomy and distraction had osteolysis instead of achieving bone length. Different explanation for pseudoarthrosis of distal tibia has been explained like thickening of periosteum, thickened and coherent periosteum which consists of



Figure 1 7 year old CPT case.



Figure 2 Unexplained resorption of the proximal bone at the site of osteotomy and distraction resulting in proximal pseudoarthrosis.



Figure 3 Grossly deformed tibia at the age of 17 years after early childhood surgery for CPT.

bone causing bone atrophy, fracture and pseudoarthrosis. Excision of hamartomatous tissue is considered as a part of treatment, which itself may not be surety of union and refracture after union. Pathology do not seems to be only localized to distal third of tibia as has been explained in osteolytic fibromatosis^[1]. Chao *et al* believes the periosteal cell has higher osteolytic activity than osteoblastic activity which may cause failure of bone healing and cause resorption of bone^[16].

In children below 5 years, we were able to apply only three rings, middle ring for distraction, one ring in lower fragment of tibia may cause anterior angulation of distal-most fragment as foot goes into planter flexion. We use sling for foot made by school of orthotics and prosthetics, Rawalpindi Medical University to prevent planter flexion. A K-wire from distal tibia to middle fragment from heel as a guide so that distraction fragment hit the distal fragment at an accurate position. Resection of sclerotic ends of pseudoarthrosis is usually done in treatment of CPT. Removing more tapered bone ends assume better cross sectional bone, better mechanical stress, chances of union and less chances of refracture^[17]. Choe *et al* observed narrow tibial shaft unite with wide proximal metaphysis has increased refracture regardless of age^[16]. Excision of more bone will increase bone gap more distraction will be required which itself become risk of fracture at distraction site. Excision of hamartomatous tissue and atrophic bone, compression of both fragments, causes early union and

avoid problem of distal and middle fragment anatomical reduction. One case we have kinking of the vessel causing pallorness of foot. It was luckily diagnosed on the table. So the distraction was given and vascularity was restored but edema of the foot remained for many weeks. The high incidence of refracture, low cross sectional area of the united segment with valgus deformity at ankle causing a stress riser and compliance issue with discarding brace earlier^[18]. Usually children are young with narrow medullary cavity, intramedullary device alone may not be sufficient. We have to rely on external bracing. In our series, we have not supplemented patients with intramedullary device. Once residual LLD has been restored, to achieve union of the distal fragment, we had to use a bone graft in two cases. Repeated traction is not indicated. Proximal migration of the distal fibula with valgus mal-alignment of ankle is seen 7/15 by Agashe *et al*. They suggest correction of the ankle valgus if greater than 5 degree is suggested by Inon *et al*.¹² BMP-2, BMP-7 and rh (recombinant human) BMP-2 has been used has not shown the promised clinical response, possible explanation is lack of osteoblastic differentiation in response to BMP. The fibrous hamartoma excision is considered as essential part of treatment^[19]. After meticulous excision of hamartomatous tissue and atrophic bone ends, achieving union at fracture site, correction of LLD by distraction histogenesis, union and consolidation of distraction fragment may not be predictable due to ill understood disease process. Refracture of the distraction fragment

or the fracture site make it more complicated. Good results with these patients are only 50%.

CONCLUSION

CPT is still a poorly understood disease perhaps primarily a periosteal disease with decreased osteoblastic activity and increased osteoclastic activity hence making the results unpredictable.

REFERENCES

- Paley D. Congenital pseudoarthrosis of the tibia: biological and biomechanical considerations to achieve union and prevent refracture. *J Child Orthop* 2019; **13**(2): 120-33. [DOI: 10.1302/1863-2548.13.180147]
- Vaidya SV, Aroojis A, Mehta R, Agashe MV, Dhawale A, Bansal AV, Sarathy K. Short Term Results of a New Comprehensive Protocol for the Management of Congenital Pseudoarthrosis of the Tibia. *Indian J Orthop* 2019; **53**(6): 736-44. [DOI: 10.4103/ortho.IJOrtho_155_19]; [PMID: 31673175]
- Eisenberg KA, Vuillermin CB. Management of Congenital Pseudoarthrosis of the Tibia and Fibula. *Curr Rev Musculoskelet Med* 2019; **12**(3): 356-68. [DOI: 10.1007/s12178-019-09566-2]; [PMID: 31228003]
- Lakloul MAS. Ilizarov fixator as a method of treatment of failed internal fixation of distal tibial fractures. *Egypt Orthop J* 2013; **48**: 88-94. [DOI: 10.7123/01.EOJ.0000426262.53476.61]
- Kong LD, Cheng HX, Nie T. Treat the congenital pseudoarthrosis of the Tibia with Ilizarov technology: Case report. *Medicine (Baltimore)* 2018; **97**(49): e13384. [DOI: 10.1097/MD.00000000000013384]; [PMID: 30544410]
- Singer D, Johnston CE. Congenital Pseudoarthrosis of the Tibia: Results, at Skeletal Maturity, of the Charnley-Williams Procedure. *JB JS Open Access* 2019; **4**(2): e0004. [DOI: 10.2106/JBJS.OA.19.00004]; [PMID: 31334459]
- Sahu RL. Percutaneous autogenous bone marrow injection for delayed union or non-union of long bone fractures after internal fixation. *Rev Bras Ortop* 2017; **53**(6): 668-673. [DOI: 10.1016/j.rboe.2017.09.004]; [PMID: 30377598]
- Zhu GH, Mei HB, He RG, Liu YX, Liu K, Tang J, Wu JY. Combination of intramedullary rod, wrapping bone grafting and Ilizarov's fixator for the treatment of Crawford type IV congenital pseudoarthrosis of the tibia: mid-term follow up of 56 cases. *BMC Musculoskelet Disord* 2016; **17**(1): 443. [DOI: 10.1186/s12891-016-1295-1]; [PMID: 27770774]
- Nicolaou N, Ghassemi A, Hill RA. Congenital pseudoarthrosis of the tibia: the results of an evolving protocol of management. *J Child Orthop* 2013; **7**(4): 269-76. [DOI: 10.1007/s11832-013-0499-2]; [PMID: 24432086]
- Patwa J, Patel R. A short series of congenital pseudoarthrosis tibia. *J Orthop* 2013; **10**(3): 123-32. [DOI: 10.1016/j.jor.2013.07.002]; [PMID: 24396228]
- Choi IH, Cho TJ, Moon HJ. Ilizarov treatment of congenital pseudoarthrosis of the tibia: a multi-targeted approach using the Ilizarov technique. *Clin Orthop Surg* 2011; **3**(1): 1-8. [DOI: 10.4055/cios.2011.3.1.1]; [PMID: 21369472]
- Agashe MV, Song SH, Refai MA, Park KW, Song HR. Congenital pseudoarthrosis of the tibia treated with a combination of Ilizarov's technique and intramedullary rodding. *Acta Orthop* 2012; **83**(5): 515-22. [DOI: 10.3109/17453674.2012.736170]; [PMID: 23043268]
- Liu Y, Mei H, Zhu G, Liu K, Wu J, Tang J, He R. Congenital pseudoarthrosis of the tibia in children: should we defer surgery until 3 years old? *J Pediatr Orthop B* 2018; **27**(1): 17-25. [DOI: 10.1097/BPB.0000000000000468]
- Joseph B, Somaraju VV, Shetty SK. Management of congenital pseudoarthrosis of the tibia in children under 3 years of age: effect of early surgery on union of the pseudoarthrosis and growth of the limb. *J Pediatr Orthop* 2003; **23**(6): 740-46. [DOI: 10.1097/00004694-200311000-00011]; [PMID:14581777]
- Grill F, Bollini G, Dungal P. Treatment approaches for congenital pseudoarthrosis of tibia: results of the EPOS multicenter study: European Paediatric Orthopaedic Society (EPOS). *J Pediatr Orthop B* 2000; **9**(2): 75-89. [DOI: 10.1097/01202412-200004000-00002]; [PMID:10868356]
- Cho T-J, Choi IH, Lee SM, Chung CY, Yoo WJ, Lee DY. Refracture after Ilizarov osteosynthesis in atrophic-type congenital pseudoarthrosis of the tibia. *J Bone Joint Surg Br* 2008; **90**(4): 488-93. [DOI: 10.1302/0301-620X.90B4.20153]
- Shah H, Joseph B, Nair BVS, Kotian DB, Choi IH, Richards BS, Johnston C, Madhuri V, Dobbs MB, Dahl M. What Factors Influence Union and Refracture of Congenital Pseudoarthrosis of the Tibia? A Multicenter Long-term Study. *J Pediatr Orthop* 2018; **38**(6): e332-37. [DOI: 10.1097/BPO.0000000000001172]
- Shah H, Rousset M, Canavese F. Congenital pseudoarthrosis of the tibia: Management and complications. *Indian J Orthop* 2012; **46**(6): 616-26. [DOI: 10.4103/0019-5413.104184]; [PMID: 23325962]
- Richards BS, Anderson TD. rhBMP-2 and Intramedullary Fixation in Congenital Pseudoarthrosis of the Tibia. *J Pediatr Orthop* 2018; **38**(4): 230-38. [DOI: 10.1097/BPO.0000000000000789]