

Possible Roles of the Transglutaminase-Catalyzed Reactions in the Pathogenesis of Human Neurodegenerative Diseases

Martina Iannaccone, Enrica Serrettiello, Nicola G. Gatta, Vittorio Gentile

Martina Iannaccone, Enrica Serrettiello, Nicola G. Gatta, Vittorio Gentile, Department of Biochemistry, Biophysics and General Pathology, Second University of Naples, via Costantinopoli 16, 80138 Naples, Italy

Correspondence to: Vittorio Gentile, Department of Biochemistry, Biophysics and General Pathology, Second University of Naples, via Costantinopoli 16, 80138 Naples, Italy.

Email: vittorio.gentile@unina2.it

Telephone: +39-81-5665870

Fax: +39-81-5665863

Received: May 20, 2015

Revised: July 28, 2015

Accepted: August 1, 2015

Published online: September 22, 2015

© 2015 ACT. All rights reserved.

Key words: Transglutaminases; Post-translational modifications of proteins; Protein aggregation; Neurodegenerative diseases; Transglutaminase inhibitors.

Iannaccone M, Serrettiello E, Gatta NG, Gentile V. Possible Roles of the Transglutaminase-Catalyzed Reactions in the Pathogenesis of Human Neurodegenerative Diseases. *Journal of Biochemistry and Molecular Biology Research* 2015; 1(3): 72-79 Available from: URL: <http://www.ghrnet.org/index.php/jbmb/article/view/1219>

ABSTRACT

Transglutaminases are Ca^{2+} -dependent enzymes which catalyze post-translational modifications of proteins. The main activity of these enzymes is the cross-linking of glutaminyl residues of a protein/peptide substrate to lysyl residues of a protein/peptide co-substrate. In addition to lysyl residues, other second nucleophilic co-substrates may include monoamines or polyamines (to form mono- or bi-substituted /crosslinked adducts) or $-\text{OH}$ groups (to form ester linkages). In absence of co-substrates, the nucleophile may be water, resulting in the net deamidation of the glutaminyl residue. Transglutaminase activity has been suggested to be involved in molecular mechanisms responsible for both physiological or pathological processes. In particular, transglutaminase activity has been shown to be responsible for a widespread human autoimmune disease, the Celiac Disease. Interestingly, neurodegenerative diseases, such as Alzheimer's Disease, Parkinson's Disease, supranuclear palsy, Huntington's Disease and other diseases, are characterized in part by aberrant cerebral transglutaminase activity and by increased cross-linked proteins in affected brains. This review focuses on the possible molecular mechanisms responsible for such diseases and on the possible use of transglutaminase inhibitors for patients with diseases characterized by aberrant transglutaminase activity.

THE BIOCHEMISTRY OF THE TRANSGLUTAMINASES

Transglutaminases (TGs, E.C. 2.3.2.13) are Ca^{2+} -dependent enzymes which catalyze post-translational modifications of proteins. Examples of TG-catalyzed reactions include: I) acyl transfer between the γ -carboxamide group of a protein/polypeptide glutaminyl residue and the γ -amino group of a protein/polypeptide lysyl residue; II) attachment of a polyamine to the γ -carboxamide of a glutaminyl residue; III) deamidation of the γ -carboxamide group of a protein/polypeptide glutaminyl residue (Figure 1)^[1,2]. The reactions catalyzed by TGs occur by a two-step mechanism (ping-pong type), (Figure 2). The transamidating activity of TGs is activated by the binding of Ca^{2+} , which exposes an active-site cysteine residue. This cysteine residue reacts with the γ -carboxamide group of an incoming glutaminyl residue of a protein/peptide substrate to yield a thioacyl-enzyme intermediate and ammonia, (Figure 2, Step 1). The thioacyl-enzyme intermediate then reacts with a nucleophilic primary amine substrate, resulting in the covalent attachment of the amine-containing donor to the substrate glutaminyl acceptor and regeneration of the cysteinyl residue at the active site, (Figure 2, Step 2). If the primary amine is donated by the ϵ -amino group of a lysyl residue in a protein/polypeptide, a N^{ϵ} -(γ -L-glutamyl)-L-lysine

(GGEL) isopeptide bond is formed, (Figure 1, example I). On the other hand, if a polyamine or another primary amine (e.g. histamine, serotonin and others) acts as the amine donor, a γ -glutamylpolyamine (or γ -glutamylamine) residue is formed, (Figure 1, example II). It is also possible for a polyamine to act as an N,N-bis-(γ -L-glutamyl) polyamine bridge between two glutaminyl acceptor residues either on the same protein/polypeptide or between two proteins/polypeptides^[3]. If there is no primary amine present, water may act as the attacking nucleophile, resulting in the deamidation of glutaminyl residues to glutamyl residues, (Figure 1, example III). Regarding the physiological roles played by the transglutaminase activity, recently transglutaminase-catalyzed polyamination of tubulin has been shown to stabilize axonal microtubules, suggesting an important role for these reactions also during some physiological processes, such as neurite outgrowth and axon maturation^[4]. The reactions catalyzed by TGs occur with little change in free energy and hence should theoretically be reversible. However, under physiological conditions the cross linking reactions catalyzed by TGs are usually irreversible. This irreversibility partly results from the metabolic removal of ammonia from the system and from thermodynamic considerations resulting from altered protein conformation. Some scientific reports suggest that TGs may be able to catalyze the hydrolysis of N^ε-(γ -L-glutamyl)-L-lysine cross-links (GGEL) isopeptide bonds in some soluble cross-linked proteins. Furthermore, it is likely that TGs can catalyze the exchange of polyamines onto proteins^[2]. In TG2 other catalytic activities, such as the ability to hydrolyze GTP (or ATP) into GDP (or ADP) and inorganic phosphate (Figure 1, example IV), a protein disulfide isomerase activity (Figure 1, example V), and a kinase activity which phosphorylates histones, retinoblastoma (RB) and P53 (Figure 1, example VI), are present, while only some of these activities have been identified also in other TGs^[5-8].

Numerous experimental evidences indicate that some TGs are multifunctional proteins with distinct and regulated enzymatic activities. In fact, under physiological conditions, the transamidation

activity of TGs is latent^[10], while other activities, recently identified, could be present. For example, in some physiological states, when the concentration of Ca^{2+} increases, the crosslinking activity of TGs may contribute to important biological processes. As previously described, one of the most intriguing properties of some TGs, such as TG2, is the ability to bind and hydrolyze GTP and, furthermore to bind to GTP and Ca^{2+} . GTP and Ca^{2+} regulate its enzymatic activities, including protein cross-linking, in a reciprocal manner: the binding of Ca^{2+} inhibits GTP-binding and GTP-binding inhibits the transglutaminase cross-linking activity of the TG2^[5]. Interestingly, TG2 shows no sequence homology with heterotrimeric or low-molecular-weight G-proteins, but there is evidence that TG2 (TG2/Gh α) is involved in signal transduction, and, therefore, TG2/Gh α should also be classified as a large molecular weight G-protein. Other studies, along with ours, showed that TG2/Gh α can mediate the activation of phospholipase C (PLC) by the α_{1b} -adrenergic receptor^[10] and can modulate adenylyl cyclase activity^[11]. TG2/Gh α can also mediate the activation of the $\delta 1$ isoform of PLC and of maxi-K channels^[12]. Interestingly, the signaling function of TG2/Gh α is preserved even with the mutagenic inactivation of its crosslinking activity by the mutation of the active site cysteine residue^[13].

THE MOLECULAR BIOLOGY OF THE TRANS-GLUTAMINASES

To date at least eight different TGs, distributed in the human body, have been identified (Table 1)^[14-19]. Complex gene expression mechanisms regulate the physiological roles that these enzymes play in both the intracellular and extracellular compartments. In the Nervous System, for example, several forms of TGs are simultaneously expressed^[20-22]. Moreover, several alternative splice variants of TGs, mostly in the 3'-end region, have been identified in these last years^[23]. Interestingly, some of them are differently expressed in human pathologies, such as Alzheimer's Disease (AD)





Figure 1 Examples of reactions catalyzed by TG: I) acyl transfer between the γ -carboxamide group of a protein/polypeptide glutaminy residue and the ε -amino group of a protein/polypeptide lysyl residue; II) attachment of a polyamine to the γ -carboxamide of a glutaminy residue; III) deamidation of the γ -carboxamide group of a protein/polypeptide glutaminy residue; IV) GTPase activity; V) protein disulfide isomerase activity; VI) protein kinase activity.

[24]. On the basis of their ubiquitous expression and their biological roles, we may speculate that the absence of these enzymes would be lethal. However, this does not always seem to be the case, since, for example, null mutants of the TG2 are usually phenotypically normal at birth^[12,25,26]. This result may be explained by the multiple expressions of other TG genes that may substitute the TG2 missing isoform, although other TG isoform mutations have been associated to severe phenotypes, such as lamellar ichthyosis for TG1 isoform mutations. Bioinformatic studies have shown that the primary structures of human TGs share some identities in only few regions, such as the active site and the calcium binding regions. However, high sequence conservation and, therefore, a high degree of preservation of secondary structure among TG2, TG3 and FXIIIa indicate that these TGs all share four-domain tertiary structures which could be similar to those of other TGs^[27].

Step 1 :



Step 2 :




Figure. 2 Schematic representation of a two step transglutaminase reaction. Step 1: In the presence of Ca^{2+} , the active-site cysteine residue reacts with the γ -carboxamide group of an incoming glutaminyl residue of a protein/peptide substrate to yield a thioacyl-enzyme intermediate and ammonia. Step 2: The thioacyl-enzyme intermediate reacts with a nucleophilic primary amine substrate, resulting in the covalent attachment of the amine-containing donor to the substrate glutaminyl acceptor and regeneration of the cysteinyl residue at the active site. If the primary amine is donated by the ϵ -amino group of a lysyl residue in a protein/polypeptide, a N^{ϵ} -(γ -L-glutamyl)-L-lysine (GGEL) isopeptide bond is formed.

POSSIBLE ROLES OF THE TRANSGLUTAMINASES IN NEURODEGENERATIVE DISEASES

Numerous scientific reports suggest that the transglutaminase activity is involved in the pathogenesis of neurodegenerative diseases, but, to date, however, also controversial experimental findings about the role of the TGs enzymes in these diseases have been obtained^[28-30]. Protein aggregates in affected brain regions are histopathological hallmarks of many neurodegenerative diseases^[31]. More than 20 years ago Selkoe *et al.*^[32] suggested that TG activity might contribute to the formation of protein aggregates in AD brain. In support of this hypothesis, tau protein has been shown to be an excellent *in vitro* substrate of TGs^[33,34] and GGEL cross-links have been found in the neurofibrillary tangles and paired helical filaments of AD brains^[35]. Interestingly, a recent work showed the presence of bis γ -glutamyl putrescine in human CSF, which was increased in Huntington's Disease (HD) CSF^[36]. This is an important evidence that protein/peptides crosslinking by polyamines does indeed occur in brain, and that this is increased in HD brain. TG activity has been shown to induce also amyloid β -protein oligomerization^[37] and aggregation at physiologic levels^[38]. By these molecular mechanisms, TGs could contribute to AD symptoms and progression^[38]. Moreover, there is evidence that TGs also contribute to the formation of proteinaceous deposits in Parkinson's Disease (PD)^[39,40], in supranuclear palsy^[41,42] and in HD, a neurodegenerative disease caused by a CAG expansion in the affected gene^[43]. For example, expanded polyglutamine domains have been reported to be substrates of TG2^[44-46] and therefore aberrant TG activity could contribute to CAG-expansion diseases, including HD (Figure 3). However, although all these studies suggest the possible involvement of the TGs in the formation of deposits of protein aggregates in neurodegenerative diseases, they do not indicate whether aberrant TG activity per se directly determines the disease progression. For example, several experimental findings reported that TG2 activity *in vitro* leads to the formation of soluble aggregates of α -synuclein^[47] or polyQ proteins^[48,49]. To date, as previously reported, at least ten human CAG-expansion diseases have been described (Table 2)^[50-59] and in at least eight of them their neuropathology is caused by the expansion in the number of residues in the polyglutamine domain to a value beyond 35-40. Remarkably, the mutated proteins have no obvious similarities except for the expanded polyglutamine domain. In fact, in all cases except SCA 12, the mutation occurs in the coding region of the gene. However, in SCA12, the CAG triplet expansion occurs in the untranslated region at the 5' end of the PPP2R2B gene. It has been proposed that the toxicity results from overexpression of the brain specific regulatory subunit of protein phosphatase PP2A^[56]. Most of the mutated proteins are widely expressed both within the brain and elsewhere in the body. A major challenge then is to understand why the brain is primarily affected and why different regions within the brain are affected in the different CAG-expansion diseases, i.e., what accounts for the neurotoxic gain of function of each protein and for a selective

Table 1 TGs and their physiological roles when known.

TG	Physiological role	Gene map location	Reference
Factor XIIIa	Blood clotting	6p24-25	[14]
TG 1 (Keratinocyte TG, kTG)	Skin differentiation	14q11.2	[15]
TG 2 (Tissue TG, tTG, cTG)	Apoptosis, cell adhesion, signal transduction	20q11-12	[16]
TG 3 (Epidermal TG, eTG)	Hair follicle differentiation	20p11.2	[17]
TG 4 (Prostate TG, pTG)	Suppression of sperm immunogenicity	3q21-2	[18]
TG 5 (TG X)	Epidermal differentiation	15q15.2	[19]
TG 6 (TG Y)	Central Nervous System development	20p13	[19]
TG 7 (TG Z)	Unknown function	15q15.2	[19]

vulnerability of each cell type. Possibly, the selective vulnerability^[60] may be explained in part by the susceptibility of the expanded polyglutamine domains in the various CAG-expansion diseases to act as cosubstrates for a brain TG (Figure 4). To strengthen the possible central role of the TGs in neurodegenerative diseases, a study by Hadjivassiliou *et al*^[61] showed that anti-TG2 IgA antibodies are present in the gut and brain of patients with gluten ataxia, a non-genetic sporadic cerebellar ataxia, but not in ataxia control patients. Recently, anti-TG2, -TG3 and -TG6 antibodies have been found in sera from CD patients, suggesting a possible involvement also of other TGs in the pathogenesis of dermatitis herpetiformis and gluten ataxia, two frequent extra intestinal manifestations of gluten sensitivity^[62,63].

In support to the hypothesis of the toxic effect of TG activity in other neurodegenerative diseases, such as Alzheimer's disease and Parkinson's Disease, TG activity has been shown to induce amyloid beta-protein and α -synuclein oligomerization and aggregation at physiologic levels^[64-66]. In fact, TG activity induces protofibril-like amyloid beta-protein assemblies that are protease-resistant and inhibit long-term potentiation^[38]. Therefore, by these molecular mechanisms, TG activity could also contribute to Alzheimer's disease symptoms and progression. Finally, a recent work by Basso *et al*^[67] found




Figure 3 Possible physiopathological effects of the mutated huntingtin. Some of the physiopathological roles of mutated huntingtin, including the formation of nuclear inclusions, have been described in the figure.

that in addition to TG2, TG1 gene expression level is significantly induced following stroke *in vivo* or due to oxidative stress *in vitro*. Moreover, structurally diverse inhibitors, used at concentrations that inhibit TG1 and TG2 simultaneously, are neuroprotective. Together,




Figure 4 Possible mechanisms responsible for protein aggregate formation catalyzed by TGs. Transglutaminase activity could produce insoluble aggregates both by the formation of N ϵ -(γ -L-glutamyl)-L-lysine (GGEL) isopeptide bonds (left side of the figure) and by the formation of N,N-bis-(γ -L-glutamyl)polyamine bridges (right side of the figure) in the mutated huntingtin.

Table 2 List of polyglutamine (CAG-expansion) diseases.

Disease	Sites of neuropathology	CAG triplet number Normal	Disease	Gene product (Intracellular localization of protein deposits)	Ref.
Corea Major or Huntington's Disease (HD)	Striatum (medium spiny neurons) and cortex in late stage	6-39	36-121	Huntingtin (n,c)	[50]
Spinocerebellar Ataxia Type 1 (SCA1)	Cerebellar cortex (Purkinje cells), dentate nucleus and brain stem	6-39	40-81	Ataxin-1 (n,c)	[51]
Spinocerebellar Ataxia Type 2 (SCA2)	Cerebellum, pontine nuclei, substantia nigra	15-29	35-64	Ataxin-2 (c)	[52]
Spinocerebellar Ataxia Type 3 (SCA3) or Machado-Joseph disease (MJD)	Substantia nigra, globus pallidus, pontine nucleus, cerebellar cortex	13-42	61-84	Ataxin-3 (c)	[53]
Spinocerebellar Ataxia Type 6 (SCA6)	Cerebellar and mild brainstem atrophy	4-18	21-30	Calcium channel Subunit (a1A) (m)	[54]
Spinocerebellar Ataxia Type 7 (SCA7)	Photoreceptor and bipolar cells, cerebellar cortex, brainstem	7-17	37-130	Ataxin-7 (n)	[55]
Spinocerebellar Ataxia Type 12 (SCA12)	Cortical, cerebellar atrophy	7-32	41-78	Brain specific regulatory subunit of protein phosphatase PP2A (?)	[56]
Spinocerebellar Ataxia Type 17 (SCA17)	Gliosis and neuronal loss in the Purkinje cell layer	29-42	46-63	TATA-binding protein (TBP) (n)	[57]
Spinobulbar Muscular Atrophy (SBMA) or Kennedy Disease	Motor neurons (anterior horn cells, bulbar neurons) and dorsal root ganglia	11-34	40-62	Androgen receptor (n, c)	[58]
Dentatorubral-pallidoluysian Atrophy (DRPLA)	Globus pallidus, dentato-rubral and subthalamic nucleus	7-35	49-88	Atrophin (n, c)	[59]

c: cytosol; m: membrane; n: nucleus.

these last studies suggested that multiple TG isoforms, not only TG2, participate in oxidative stress-induced cell death signalling, and that isoform nonselective inhibitors of TG will be most efficacious in combating oxidative death in neurological disorders. These are interesting and worthwhile studies, suggesting that multiple TG isoforms can participate in neuronal death processes. Therefore, all these studies suggest that the involvement of brain TGs could represent a common denominator in several neurological diseases, which can lead to the determination of pathophysiological consequences through different molecular mechanisms.

TRANSGLUTAMINASE INHIBITION AS POSSIBLE THERAPEUTICAL APPROACH

In consideration to the fact that up to now there have been no long-term effective treatments for the human neurodegenerative diseases previously reported, then the possibility that selective TG inhibitors may be of clinical benefit has been seriously considered. In this respect, some encouraging results have been obtained with TG inhibitors in preliminary studies with different biological models of CAG-expansion diseases. For example, cystamine (Figure 5), is a potent *in vitro* inhibitor of enzymes that require an unmodified cysteine at the active site^[68]. Inasmuch as TGs contain a crucial active-site cysteine, cystamine has the potential to inhibit these enzymes by a sulfide-disulfide interchange reaction. A sulfide-disulfide interchange reaction results in the formation of cysteamine and a cysteamine-cysteine mixed disulfide residue at the active site. Recent studies have shown that cystamine decreases the number of protein inclusions in transfected cells expressing the atrophin (DRPLA) protein containing a pathological-length polyglutamine domain^[69]. In other studies, cystamine administration to HD-transgenic mice resulted in an increase in life expectancy and amelioration of neurological symptoms^[70,71]. Neuronal inclusions were decreased in one of these studies^[71]. Although all these scientific reports seem to support the hypothesis of a direct role of transglutaminase activity in the pathogenesis of the polyglutamine diseases, cystamine is also found to act in the HD-transgenic mice by mechanisms other than the inhibition of TGs, such as the inhibition of caspases^[72], suggesting that this compound can have an additive effect in the therapy of HD. Currently, cysteamine is already in phase I studies in humans with HD^[73], but several side effects, such as nausea, motor impairment and dosing schedule have been reported as reasons for non-adherence during phase II studies in human patients affected by cystinosis^[74,75]. Another critical problem in the use of TG inhibitors in treating neurological diseases relates to the fact that, as previously reported, the human brain contains at least four TGs, including TG1, 2, 3^[22] and TG6^[76], and a strong non-selective inhibitor of TGs might also inhibit plasma Factor XIIIa, causing a bleeding disorder. Therefore, from a number of standpoints it would seem that a selective inhibitor, which discriminates between TGs, would be preferable to an indiscriminate TG inhibitor. In fact, although most of the TG activity in mouse brain, at least as assessed by an assay that measures the incorporation of radioactive putrescine (amine donor) into N,N-dimethyl casein (amine acceptor), seems to be due to TG2^[77], no conclusive data have been obtained by TG2




Figure 5 Chemical structure of cystamine.

gene knock-out experiments about the involvement of this TG in the development of the symptoms in HD-transgenic mice^[26,78,79]. Moreover, a recent scientific report showed that cystamine reduces aggregate formation in a mouse model of oculopharyngeal muscular dystrophy (OMPD), in which also the TG2 knockdown is capable of suppressing the aggregation and the toxicity of the mutant protein PABPN1^[80], suggesting this compound as a possible therapeutic for OMPD.

CONCLUSIONS

In conclusion, numerous scientific reports have implicated aberrant TG activity in neurodegenerative diseases, but still today we are looking for experimental findings which could definitely confirm the direct involvement of TGs in the pathogenetic mechanisms responsible for these diseases. However, as result of the putative role of specific TG isoforms, such as TG2, in some human diseases, there is a considerable interest in developing inhibitors of these enzymes. Of those currently available, cystamine is the most commonly used experimentally to inhibit TG2 activity. In addition to cystamine, several types of TG2 inhibitors have been developed up to now^[81]. Interestingly, some of these inhibitors have shown promising results in experimental diabetic models^[82]. Therefore, the use of these inhibitors of TGs could be then useful also for other clinical approaches. To minimize the possible side effects, however, more selective inhibitors of the TGs should be required in the future. Progress in this area of research could be achieved also through pharmaco-genetic techniques.

ACKNOWLEDGEMENTS

This work is supported by the Italian Education Department and the Regione Campania (L.R. n.5 del 28.03.2002, finanziamento 2008).

CONFLICT OF INTERESTS

The Authors have no conflicts of interest to declare.

REFERENCES

- 1 Folk JE. Mechanism and basis for specificity of transglutaminase-catalyzed ϵ -(γ -glutamyl)lysine bond formation. *Adv Enzymol Relat Areas Mol Biol* 1983; 54: 1-56.
- 2 Lorand L, Conrad S M. Transglutaminases. *Mol Cell Biochem* 1984; 58: 9-35.
- 3 Piacentini M, Martinet N, Beninati S, Folk J E. Free and protein conjugated-polyamines in mouse epidermal cells. Effect of high calcium and retinoic acid. *J Biol Chem* 1988; 263: 3790-3794.
- 4 Song Y, Kirkpatrick LL, Schilling AB, Helseth DL, Chabot N, Keillor JW, Johnson GV, Brady ST. Transglutaminase and polyamination of tubulin: posttranslational modification for stabilizing axonal microtubules. *Neuron* 2013; 78: 109-123.
- 5 Achyuthan KE, Greenberg CS. Identification of a guanosine triphosphate-binding site on guinea pig liver transglutaminase. Role of GTP and calcium ions in modulating activity. *J Biol Chem* 1987; 262: 1901-1906.
- 6 Hasegawa G, Suwa M, Ichikawa Y, Ohtsuka T, Kumagai S, Kikuchi M, Sato Y, Saito Y. A novel function of tissue-type transglutaminase: protein disulfide isomerase. *Biochem J* 2003; 373: 793-803.
- 7 Lahav J, Karniel E, Bagoly Z, Sheptovitsky V, Dardik R, Inbal A. Coagulation factor XIII serves as protein disulfide isomerase. *Thromb Haemost* 2009; 101: 840-844.
- 8 Iismaa SE, Mearns BM, Lorand L, Graham RM. Transglutaminas-

- es and disease: lessons from genetically engineered mouse models and inherited disorders. *Physiol Rev* 2009; 89: 991-1023.
- 9 Smethurst PA, Griffin M. Measurement of tissue transglutaminase activity in a permeabilized cell system: its regulation by calcium and nucleotides. *Biochem J* 1996; 313: 803-808.
 - 10 Nakaoka H, Perez DM, Baek KJ, Das T, Husain A, Misono K, Im MJ, Graham RM. Gh: a GTP-binding protein with transglutaminase activity and receptor signalling function. *Science* 1994; 264: 1593-1596.
 - 11 Gentile V, Porta R, Chiosi E, Spina A, Caputo I, Valente F, Pezone R, Davies PJA, Illiano G. Tissue transglutaminase and adenylate cyclase interactions in Balb-C 3T3 fibroblast membranes. *Biochim Biophys Acta* 1997; 1357: 115-122.
 - 12 Nanda N, Iismaa SE, Owens WA, Husain A, Mackay F, Graham RM. Targeted inactivation of Gh/tissue transglutaminase II. *J Biol Chem* 2001; 276: 20673-20678.
 - 13 Mian S, El Alaoui S, Lawry J, Gentile V, Davies PJA, Griffin M. The importance of the GTP binding protein tissue transglutaminase in the regulation of cell cycle progression. *FEBS Letters* 1995; 370: 27-31.
 - 14 Olaisen B, Gedde-Dahl TJR, Teisberg P, Thorsby E, Siverts A, Jonassen R, Wilhelmy MC. A structural locus for coagulation factor XIIIa (F13A) is located distal to the HLA region on chromosome 6p in man. *Am J Hum Genet* 1985; 37: 215-220.
 - 15 Yamanishi K, Inazawa J, Liew F-M, Nonomura K, Ariyama T, Yasuno H, Abe T, Doi H, Hirano J, Fukushima S. Structure of the gene for human transglutaminase 1. *J Biol Chem* 1992; 267: 17858-17863.
 - 16 Gentile V, Davies PJA, Baldini A. The human tissue transglutaminase gene maps on chromosome 20q12 by in situ fluorescence hybridization. *Genomics* 1994; 20: 295-297.
 - 17 Wang M, Kim IG, Steinert PM, McBride OW. Assignment of the human transglutaminase 2 (TGM2) and transglutaminase 3 (TGM3) genes to chromosome 20q11.2. *Genomics* 1994; 23: 721-722.
 - 18 Gentile V, Grant F, Porta R, Baldini A. Human prostate transglutaminase is localized on chromosome 3p21.33-p22 by in situ fluorescence hybridization. *Genomics* 1995; 27: 219-220.
 - 19 Grenard P, Bates MK, Aeschlimann D. Evolution of transglutaminase genes: identification of a transglutaminases gene cluster on human chromosome 15q. Structure of the gene encoding transglutaminase X and a novel gene family member, transglutaminase Z. *J Biol Chem* 2001; 276: 33066-33078.
 - 20 Thomas H, Beck K, Adamczyk M, Aeschlimann P, Langley M, Oita RC, Thiebach L, Hils M, Aeschlimann D. Transglutaminase 6: a protein associated with central nervous system development and motor function. *Amino Acids* 2013; 44: 161-177.
 - 21 Bailey CDC, Johnson GVW. Developmental regulation of tissue transglutaminase in the mouse forebrain. *J Neurochem* 2004; 91: 1369-1379.
 - 22 Kim S-Y, Grant P, Lee JHC, Pant HC, Steinert PM. Differential expression of multiple transglutaminases in human brain. Increased expression and cross-linking by transglutaminase 1 and 2 in Alzheimer's disease. *J Biol Chem* 1999; 274: 30715-30721.
 - 23 Iannaccone M, Giuberti G, De Vivo G, Caraglia M, Gentile V. Identification of a FXIIIa variant in human neuroblastoma cell lines. *Int J Biochem Mol Biol* 2013; 4: 102-107.
 - 24 Citron BA, Santa Cruz KS, Davies PJ, Festoff BW. Intron-exon swapping of transglutaminase mRNA and neuronal tau aggregation in Alzheimer's disease. *J Biol Chem* 2001; 276: 3295-3301.
 - 25 De Laurenzi V, Melino G. Gene disruption of tissue transglutaminase. *Mol Cell Biol* 2001; 21: 148-155.
 - 26 Mastroberardino PG, Iannicola C, Nardacci R, Bernassola F, De Laurenzi V, Melino G, Moreno S, Pavone F, Oliverio S, Fesus L, Piacentini M. 'Tissue' transglutaminase ablation reduces neuronal death and prolongs survival in a mouse model of Huntington's disease. *Cell Death and Differentiation* 2002; 9: 873-880.
 - 27 Lorand L, Graham RM. Transglutaminases: crosslinking enzymes with pleiotropic functions. *Nature Mol Cell Biol* 2003; 4: 140-156.
 - 28 Wolf J, Jäger C, Lachmann I, Schönknecht P, Morawski M, Arendt T, Mothes T. Tissue transglutaminase is not a biochemical marker for Alzheimer's disease. *Neurobiol Aging* 2013; 34: 2495-2498.
 - 29 Wilhelmus MMM, Drukarch B. Tissue transglutaminase is a biochemical marker for Alzheimer's disease. *Neurobiol Aging* 2014; 35: e3-e4.
 - 30 Wolf J, Jäger C, Morawski M, Lachmann I, Schönknecht P, Mothes T, Arendt T. Tissue transglutaminase in Alzheimer's disease - facts and fiction: a reply to "Tissue transglutaminase is a biochemical marker for Alzheimer's disease". *Neurobiol Aging* 2014; 35: e5-e9.
 - 31 Adams RD, Victor M. Principles of Neurology. McGraw-Hill, Inc. Ed. 1993.
 - 32 Selkoe DJ, Abraham C, Ihara Y. Alzheimer's disease: insolubility of partially purified paired helical filaments in sodium dodecyl sulfate and urea. *Proc Natl Acad Sci U S A* 1982; 79: 6070-6074.
 - 33 Grierson AJ, Johnson GV, Miller CC. Three different human τ isoforms and rat neurofilament light, middle and heavy chain proteins are cellular substrates for transglutaminase. *Neurosci Lett* 2001; 298: 9-12.
 - 34 Singer SM, Zainelli GM, Norlund MA, Lee JM, Muma NA. Transglutaminase bonds in neurofibrillary tangles and paired helical filament τ early in Alzheimer's disease. *Neurochem Int* 2002; 40: 17-30.
 - 35 Halverson RA, Lewis J, Frausto S, Hutton M, Muma NA. Tau protein is cross-linked by transglutaminase in P301L tau transgenic mice. *J Neurosci* 2005; 25(5): 1226-33.
 - 36 Jeitner TM, Matson WR, Folk JE, Blass JP, Cooper AJL. Increased levels of γ -glutamylamines in Huntington disease CSF. *J Neurochem* 2008; 106: 37-44.
 - 37 Dudek SM, Johnson GV. Transglutaminase facilitates the formation of polymers of the beta-amyloid peptide. *Brain Res* 1994; 651(1-2):129-33.
 - 38 Hartley DM, Zhao C, Speier AC, Woodard GA, Li S, Li Z, Walz T. Transglutaminase induces protofibril-like amyloid β -protein assemblies that are protease-resistant and inhibit long-term potentiation. *J Biol Chem* 2008; 283: 16790-16800.
 - 39 Citron BA, Suo Z, SantaCruz K, Davies PJ, Qin F, Festoff BW. Protein crosslinking, tissue transglutaminase, alternative splicing and neurodegeneration. *Neurochem Int* 2002; 40: 69-78.
 - 40 Junn E, Ronchetti RD, Quezado MM, Kim SY, Mouradian MM. Tissue transglutaminase-induced aggregation of α -synuclein: Implications for Lewy body formation in Parkinson's disease and dementia with Lewy bodies. *Proc Natl Acad Sci USA* 2003; 100: 2047-2052.
 - 41 Zemaitaitis MO, Lee JM, Troncoso JC, Muma NA. Transglutaminase-induced cross-linking of τ proteins in progressive supranuclear palsy. *J Neuropathol Exp Neurol* 2000; 59: 983-989.
 - 42 Zemaitaitis MO, Kim SY, Halverson RA, Troncoso JC, Lee JM, Muma NA. Transglutaminase activity, protein, and mRNA expression are increased in progressive supranuclear palsy. *J Neuropathol Exp Neurol* 2003; 62: 173-184.
 - 43 Iuchi S, Hoffner G, Verbeke P, Djian P, Green H. Oligomeric and polymeric aggregates formed by proteins containing expanded polyglutamine. *Proc Natl Acad Sci U S A* 2003; 100: 2409-2414.
 - 44 Gentile V, Sepe C, Calvani M, Melone MAB, Cotrufo R, Cooper AJL, Blass JP and Peluso G. Tissue transglutaminase-catalyzed formation of high-molecular-weight aggregates in vitro is favored with long polyglutamine domains: a possible mechanism contributing to CAG-triplet diseases. *Arch Biochem Biophys* 1998; 352: 314-321.
 - 45 Kahlem P, Green H, Djian P. Transglutaminase action imitates Huntington's disease: selective polymerization of huntingtin containing expanded polyglutamine. *Mol Cell* 1998; 1: 595-601.
 - 46 Karpuj MV, Garren H, Slunt H, Price DL, Gusella J, Becher MW, Steinman L. Transglutaminase aggregates huntingtin into non-

- amyloidogenic polymers, and its enzymatic activity increases in Huntington's disease brain nuclei. *Proc Natl Acad Sci U S A* 1999; 96: 7388-7393.
- 47 Segers-Nolten IM, Wilhelmus MM, Veldhuis G, van Rooijen BD, Drukarch B, Subramaniam V. Tissue transglutaminase modulates α -synuclein oligomerization. *Protein Sci* 2008; 17: 1395-1402.
- 48 Lai T-S, Tucker T, Burke JR, Strittmatter WJ, Greenberg CS. Effect of tissue transglutaminase on the solubility of proteins containing expanded polyglutamine repeats. *J Neurochem* 2004; 88: 1253-1260.
- 49 Konno T, Mori T, Shimizu H, Oiki S, Ikura K. Paradoxical inhibition of protein aggregation and precipitation by transglutaminase-catalyzed intermolecular cross-linking. *J Biol Chem* 2005; 280: 17520-17525.
- 50 The Huntington's Disease Collaborative Research Group. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosome. *Cell* 1993; 72: 971-983.
- 51 Banfi S, Chung MY, Kwiatkowski TJ Jr, Ranum LP, McCall AE, Chinault AC, Orr HT, Zoghbi HY. Mapping and cloning of the critical region for the spinocerebellar ataxia type 1 gene (SCA1) in a yeast artificial chromosome contig spanning 1.2 Mb. *Genomics* 1993; 18: 627-635.
- 52 Sanpei K, Takano H, Igarashi S, Sato T, Oyake M, Sasaki H, Wakisaka A, Tashiro K, Ishida Y, Ikeuchi T, Koide R, Saito M, Sato A, Tanaka T, Hanyu S, Takiyama Y, Nishizawa M, Shimizu N, Nomura Y, Segawa M, Iwabuchi K, Eguchi I, Tanaka H, Takahashi H, Tsuji S. Identification of the spinocerebellar ataxia type 2 gene using a direct identification of repeat expansion and cloning technique, DIRECT. *Nat Genet* 1996; 14: 277-284.
- 53 Pujana MA, Volpini V, Estivill X. Large CAG/CTG repeat templates produced by PCR, usefulness for the DIRECT method of cloning genes with CAG/CTG repeat expansions. *Nucleic Acids Res* 1998; 1: 1352-1353.
- 54 Fletcher CF, Lutz CM, O'Sullivan TN, Shaughnessy JD Jr, Hawkes R, Frankel WN, Copeland NG, Jenkins NA. Absence epilepsy in tottering mutant mice is associated with calcium channel defects. *Cell* 1996; 87: 607-617.
- 55 Vincent JB, Neves-Pereira ML, Paterson AD, Yamamoto E, Parikh SV, Macciardi F, Gurling HM, Potkin SG, Pato CN, Macedo A, Kovacs M, Davies M, Lieberman JA, Meltzer HY, Petronis A, Kennedy JL. An unstable trinucleotide-repeat region on chromosome 13 implicated in spinocerebellar ataxia: a common expansion locus. *Am J Hum Genet* 2000; 66: 819-829.
- 56 Holmes SE, O'Hearn E, Margolis RL. Why is SCA12 different from other SCAs? *Cytogenet Genome Res* 2003; 100: 189-197.
- 57 Imbert G, Trotter Y, Beckmann J, Mandel JL. The gene for the TATA binding protein (TBP) that contains a highly polymorphic protein coding CAG repeat maps to 6q27. *Genomics* 1994; 21: 667-668.
- 58 La Spada AR, Wilson EM, Lubahn DB, Harding AE, Fischbeck KH. Androgen receptor gene mutations in X-linked spinal and bulbar muscular atrophy. *Nature* 1991; 352: 77-79.
- 59 Onodera O, Oyake M, Takano H, Ikeuchi T, Igarashi S, Tsuji S. Molecular cloning of a full-length cDNA for dentatorubral-pallidoluysian atrophy and regional expressions of the expanded alleles in the CNS. *Am J Hum Genet* 1995; 57: 1050-1060.
- 60 Cooper AJL, Sheu K-FR, Burke JR, Strittmatter WJ, Gentile V, Peluso G, and Blass JP. Pathogenesis of inclusion bodies in (CAG) n/Qn-expansion diseases with special reference to the role of tissue transglutaminase and to selective vulnerability. *J Neurochem* 1999; 72: 889-899.
- 61 Hadjivassiliou M, Maki M, Sanders DS, Williamson CA, Grunewald RA, Woodroffe NM, Korponay-Szabó IR. Autoantibody targeting of brain and intestinal transglutaminase in gluten ataxia. *Neurology* 2006; 66: 373-377.
- 62 Boscolo S, Lorenzon A, Sblattero D, Florian F, Stebel M, Marzari R, Not T, Aeschlimann D, Ventura A, Hadjivassiliou M, Tongiorgi E. Anti transglutaminase antibodies cause ataxia in mice. *PLoS One* 2010 Mar 15; 5 (3): e9698.
- 63 Stammaes J, Dorum S, Fleckenstein B, Aeschlimann D, Sollid LM. Gluten T cell epitope targeting by TG3 and TG6; implications for dermatitis herpetiformis and gluten ataxia. *Amino Acids* 2010; 39: 1183-1191.
- 64 Wakshlag JJ, Antonyak MA, Boehm JE, Boehm K, Cerione RA. Effects of tissue transglutaminase on beta-amyloid 1-42-induced apoptosis. *Protein J* 2006; 25: 83-94.
- 65 Lee JH, Jeong J, Jeong EM, Cho SY, Kang JW, Lim J, Heo J, Kang H, Kim IG, Shin DM. Endoplasmic reticulum stress activates transglutaminase 2 leading to protein aggregation. *Int J Mol Med* 2014; 33: 849-55.
- 66 Grosso H, Woo JM, Lee KW, Im JY, Masliah E, Junn E, Mouradian MM. Transglutaminase 2 exacerbates α -synuclein toxicity in mice and yeast. *FASEB J* 2014; 28(10): 4280-91.
- 67 Basso M, Berlin J, Xia L, Sleiman SF, Ko B, Haskew-Layton R, Kim E, Antonyak MA, Cerione RA, Iismaa SE, Willis D, Cho S, Ratan RR. Transglutaminase inhibition protects against oxidative stress-induced neuronal death downstream of pathological ERK activation. *Journal of Neurosciences* 2012; 39: 6561-6569.
- 68 Griffith OW, Larsson A, Meister A. Inhibition of γ -glutamylcysteine synthetase by cystamine: an approach to a therapy of 5-oxoprolinuria (pyroglutamic aciduria). *Biochem Biophys Res Commun* 1977; 79: 919-925.
- 69 Igarashi S, Koide R, Shimohata T, Yamada M, Hayashi Y, Takano H, Date H, Oyake M, Sato T, Sato A, Egawa S, Ikeuchi T, Tanaka H, Nakano R, Tanaka K, Hozumi I, Inuzuka T, Takahashi H, Tsuji S. Suppression of aggregate formation and apoptosis by transglutaminase inhibitors in cells expressing truncated DRPLA protein with an expanded polyglutamine stretch. *Nat Genet* 1998; 18: 111-117.
- 70 Karpuj MV, Becher MW, Springer JE, Chabas D, Youssef S, Pedotti R, Mitchell D, Steinman L. Prolonged survival and decreased abnormal movements in transgenic model of Huntington disease, with administration of the transglutaminase inhibitor cystamine. *Nature Medicine* 2002; 8: 143-149.
- 71 Dedeoglu A, Kubilus JK, Jeitner TM, Matson SA, Bogdanov M, Kowall NW, Matson WR, Cooper AJ, Ratan RR, Beal MF, Hersch SM, Ferrante RJ. Therapeutic effects of cystamine in a murine model of Huntington's disease. *J Neurosci* 2002; 22: 8942-8950.
- 72 Lesort M, Lee M, Tucholski J, Johnson GVV. Cystamine inhibits caspase activity. Implications for the treatment of polyglutamine disorders. *J Biol Chem* 2003; 278: 3825-3830.
- 73 Dubinsky R, Gray C. CYTE-I-HD: Phase I Dose Finding and Tolerability Study of Cystamine (Cystagon) in Huntington's Disease. *Movement Disorders* 2006; 21: 530-533.
- 74 Langman CB, Greenbaum LA, Sarwal M, Grimm P, Niaudet P, Deschènes G, Cornelissen E, Morin D, Cochat P, Matossian D, Gaillard S, Bagger MJ, Rioux P. A randomized controlled cross-over trial with delayed-release cystamine bitartrate in nephropathic cystinosis: effectiveness on white blood cell cystine levels and comparison of safety. *Clin J Am Soc Nephrol* 2012; 7: 1112-20.
- 75 Besouw M, Masereeuw R, van den Heuvel L, Levchenko E. Cystamine: an old drug with new potential. *Drug Discovery Today* 2013; 18: 785-792.
- 76 Hadjivassiliou M, Aeschlimann P, Strigun A, Sanders DS, Woodroffe N, Aeschlimann D. Autoantibodies in gluten ataxia recognize a novel neuronal transglutaminase. *Ann Neurol* 2008; 64: 332-343.
- 77 Krasnikov BF, Kim SY, McConoughey SJ, Ryu H, Xu H, Stavrovskaya I, Iismaa SE, Mearns BM, Ratan RR, Blass JP, Gibson GE, Cooper AJ. Transglutaminase activity is present in highly purified nonsynaptosomal mouse brain and liver mitochondria. *Biochemistry* 2005; 44: 7830-7843.

- 78 Menalled LB, Kudwa AE, Oakeshott S, Farrar A, Paterson N, Filippov I, Miller S, Kwan M, Olsen M, Beltran J, Torello J, Fitzpatrick J, Mushlin R, Cox K, McConnell K, Mazzella M, He D, Osborne GF, Al-Nackkash R, Bates GP, Tuunanen P, Lehtimäki K, Brunner D, Ghavami A, Ramboz S, Park L, Macdonald D, Munoz-Sanjuan I, Howland D. Genetic deletion of transglutaminase 2 does not rescue the phenotypic deficits observed in R6/2 and zQ175 mouse models of Huntington's disease. *PLoS One* 2014 Jun 23; 9(6):e99520. doi: 10.1371/journal.pone.0099520. eCollection 2014.
 - 79 Bailey CD, Johnson GV. Tissue transglutaminase contributes to disease progression in the R6/2 Huntington's disease mouse model via aggregate-independent mechanisms. *J Neurochem* 2005; 92: 83-92.
 - 80 Davies JE, Rose C, Sarkar S, Rubinsztein DC. Cystamine suppresses polyalanine toxicity in a mouse model of oculopharyngeal muscular dystrophy. *Science Translational Medicine* 2010; 2: 34-40.
 - 81 Pietsch M, Wodtke R, Pietzsch J, Löser R. Tissue transglutaminase: An emerging target for therapy and imaging. *Bioorganic & Medicinal Chemistry Letters* 2013; 23: 6528–6543.
 - 82 Bhatt MP, Lim YC, Hwang J, Na S, Kim YM, Ha KS. C-peptide prevents hyperglycemia-induced endothelial apoptosis through inhibition of reactive oxygen species-mediated transglutaminase 2 activation. *Diabetes* 2013, 62: 243-253.
- Peer reviewer:** Qun Bi, Associate Prof., College of Life Sciences, Peking University, Yiheyuan Street #5, Beijing 100871, China.